

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 May 2001 (10.05.2001)

PCT

(10) International Publication Number
WO 01/32014 A2

(51) International Patent Classification⁷: A01N

(21) International Application Number: PCT/US00/30191

(22) International Filing Date:
1 November 2000 (01.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/431,705 1 November 1999 (01.11.1999) US

(71) Applicant: ORAVAX, INC. [US/US]; 38 Sidney Street,
Cambridge, MA 02139-4169 (US).

(72) Inventors: KLEANTHOUS, Harold; 89 Madison
Avenue, Newtonville, MA 02160 (US). LONDONO-AR-
CILA, Patricia; Flat B, 11 Beckwith Road, London
SE24 9LH (GB). FREEMAN, Donna; 68 Thorpe Way,
Cambridge CB5 8UB (GB).

(74) Agent: MICHAUD, Susan, M.; Clark & Elbing LLP, 176
Federal Street, Boston, MA 02110-2214 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: USE OF SALMONELLA VECTORS FOR VACCINATION AGAINST HELICOBACTER INFECTION



(57) Abstract: The invention provides a method of immunization against Helicobacter, involving mucosal administration of an attenuated Salmonella vector including a nucleic acid molecule encoding a Helicobacter antigen, and parenteral administration of a soluble Helicobacter antigen, co-administered with a suitable parenteral adjuvant. Also provided by the invention are attenuated Salmonella vectors for use in this method.

USE OF SALMONELLA VECTORS FOR VACCINATION AGAINST HELICOBACTER INFECTION

Background of the Invention

This invention relates to the use of Salmonella vectors in vaccination methods against Helicobacter infection.

Helicobacter is a genus of spiral, gram-negative bacteria that colonize the gastrointestinal tracts of mammals. Several species colonize the stomach, most notably *H. pylori*, *H. heilmanii*, *H. felis*, and *H. mustelae*. Although *H. pylori* is the species most commonly associated with human infection, *H. heilmanii* and *H. felis* have also been isolated from humans, but at lower frequencies than *H. pylori*. Helicobacter infects over 50% of adult populations in developed countries and nearly 100% in developing countries and some Pacific rim countries, making it one of the most prevalent infections worldwide.

Helicobacter is routinely recovered from gastric biopsies of humans with histological evidence of gastritis and peptic ulceration. Indeed, *H. pylori* is now recognized as an important pathogen of humans, in that the chronic gastritis it causes is a risk factor for the development of peptic ulcer diseases and gastric carcinoma. It is thus highly desirable to develop safe and effective methods for preventing and treating Helicobacter infection.

Summary of the Invention

The invention provides a method of inducing an immune response against *Helicobacter* in a mammal. This method involves mucosally (*e.g.*, orally) administering to a mammal (*e.g.*, a human) an attenuated *Salmonella* (*e.g.*, *S. typhi* (*e.g.*, CVD908-htrA or CVD908) or *S. typhimurium* (*e.g.*,

BRD509 or BRD807)) vector including a nucleic acid molecule encoding a Helicobacter antigen (e.g., a urease, a urease subunit, or an immunogenic fragment thereof), and parenterally administering to the mammal a Helicobacter antigen (e.g., a urease, a urease subunit, or an immunogenic fragment thereof), optionally, in association with an adjuvant, such as an aluminum compound (e.g., alum). The nucleic acid molecule encoding the Helicobacter antigen can be under the control of a promoter, such as an *htrA* or a *nirB* promoter. The antigen used in the mucosal administration can be different from, cross-reactive with, or, preferably, identical to the parenterally administered antigen. In a preferred embodiment, the mucosal administration primes an immune response to an antigen, and the parenteral administration boosts an immune response to the antigen. A mammal treated according to the method of the invention can be at risk of developing, but not have, a Helicobacter infection, or can have a Helicobacter infection. That is, the method can be used to prevent or to treat Helicobacter infection.

The invention also includes use of the Salmonella vectors described above in the preparation of medicaments for preventing or treating Helicobacter (e.g., *Helicobacter pylori*) infection by mucosal (e.g., oral) administration of the vectors, and parenteral administration of a Helicobacter antigen (e.g., a urease, a urease subunit, or an immunogenic fragment thereof; also see other antigens listed herein), optionally in association with an adjuvant (e.g., alum).

The invention also provides an attenuated Salmonella (e.g., *S. typhi* (e.g., CVD908-htrA or CVD908) or *S. typhimurium* (e.g., BRD509 or BRD807)) vector including a nucleic acid molecule encoding a Helicobacter antigen, e.g., a urease, a urease subunit, or an immunogenic fragment thereof, expressed as a fusion protein that can be selectively

targeted to the outer membrane or secreted from the cell. The nucleic acid molecule encoding the Helicobacter antigen can be under the control of a promoter, such as an *htrA* or a *nirB* promoter.

Other features and advantages of the invention will be apparent from the following detailed description, the drawings, and the claims.

Brief Description of the Drawings

Fig. 1 is a schematic representation of an expression plasmid (pH/NUR3) used in Salmonella immunizations.

Fig. 2A is a graph showing the urease-specific serum antibody (IgG2a) response of mice that were mucosally primed with *S. typhimurium*-vectored urease, followed by parenteral boosting with urease and alum.

Fig. 2B is a graph showing the T helper phenotype (IgG1/IgG2a ratio) of mice that were mucosally primed with *S. typhimurium*-vectored urease, followed by parenteral boosting with urease and alum.

Fig. 3A is a graph showing protection against Helicobacter infection in mice that were mucosally primed with *S. typhimurium*-vectored urease, followed by parenteral boosting with urease and alum.

Fig. 3B is a table showing protection against Helicobacter infection in mice that were mucosally primed with *S. typhimurium*-vectored urease, followed by parenteral boosting with urease and alum, as log₁₀ reduction in comparison to a no treatment control group.

Fig. 4 provides the nucleic acid sequence (SEQ ID NOs:1 and 19) and amino acid sequence (SEQ ID NOs:2-18 and 20-30) of plasmid pHUR3.

Fig. 5 is a schematic representation of some relevant features of pHUR3.

Detailed Description

This invention provides an immunization method against *Helicobacter* infection that involves: (i) mucosal administration of an attenuated *Salmonella* vector containing a nucleic acid molecule encoding a *Helicobacter* antigen, and (ii) parenteral administration of a *Helicobacter* antigen, preferably, in association with an adjuvant. The method can be used to prevent or to treat *Helicobacter* infection in a mammal, such as a human. Also, the mucosal administration can be used to prime an immune response to an antigen, and the parenteral administration can be used to boost an immune response to the antigen. The invention also provides *Salmonella* vectors for use in this method. *Salmonella* vectors, *Helicobacter* antigens, and adjuvants that can be used in the method of the invention are first described, as follows. Then, details of the immunization method of the invention, and examples of its efficacy, are provided.

Salmonella Vectors

Numerous attenuated *Salmonella* vectors that can be used in the invention are known in the art, and can be derived from species such as, for example, *S. typhi*, *S. typhimurium*, *S. enteritidis*, *S. dublin*, *S. minnesota*, and *S. choleraesuis*. The vectors can be attenuated chemically (e.g., Ty21a, Swiss Serums and Vaccines, Berna Products) or, preferably, by genetic mutagenesis (e.g., Ty800). For example, attenuation can be achieved by inactivation of key regulatory genes or genes necessary for *in vivo* survival. For example, the following genes can be inactivated: *cya*, *crp*, and *asd* (cAMP metabolism; see, e.g., Curtiss *et al.*, Vaccine 6:155-160, 1988; Nakayama *et al.*, BioTechnology 6:693, 1988; WO 92/11361), adenylate cyclase and the cAMP receptor (U.S. Patent No. 5,389,368), *cdt* (invasion of liver and spleen), *phoP/phoQ* (two component regulator; see, e.g., Fields

et al., Science 243:1059-1062, 1989; U.S. Patent No. 5,424,065), *ompR* (control of capsule and porin expression; see, *e.g.*, Dorman *et al.*, Infection and Immunity 57:2136-2140, 1989), outer membrane proteins (U.S. Patent No. 5,527,529), reverse mutants of streptomycin mutants (U.S. Patent No. 4,350,684), genes in pathogenicity islands (Shea *et al.*, Infection and Immunity 67:213-219, 1999; WO 99/37759), *SPI-2* (invasion of Peyer's patches), *Dam* (DNA methylation), *htrA* (heat shock protein; U.S. Patent No. 5,804,194), and other heat shock proteins (U.S. Patent No. 5,804,194). The vectors can also be attenuated by auxotrophic mutations, such as mutations in any of the *aroA*, *aroC*, *aroD* (aromatic compounds), *purA*, or *guaAB* (purines) genes (see, *e.g.*, U.S. Patent No. 5,770,214).

Preferably, the mutations in the Salmonella strains used in the invention are non-reverting mutations, *i.e.*, mutations that cannot be repaired in a single step. Mutations of this sort include deletions, inversions, insertions, and substitutions. Preferably, there is more than one mutation in the vector. Methods of making such mutations are well known in the art.

Specific examples of Salmonella vectors that can be used in the invention include *S. typhi* mutant strains, for example, CVD908 *S. typhi* Ty2 Δ aroC/ Δ aroD (Hone *et al.*, Vaccine 9:810-816, 1991), CVD908-*htrA* *S. typhi* Ty2 Δ aroC/ Δ aroD/ Δ htrA (Tacket *et al.*, Infection and Immunity 65:452-456, 1997), BRD1116 *S. typhi* Ty2 Δ aroA/ Δ aroC/ Δ htrA (Lowe *et al.*, Infection and Immunity 67:700-707, 1999), *S. typhi* Δ aroA/ Δ aroE (U.S. Patent No. 5,770,214; deposited at PHLS, NCTC, 61 Colindale Avenue, London NW9 5HT under Accession No. NCTC 12164), *S. typhi* Ty2 Δ aroA/ Δ aroC Km-R (U.S. Patent No. 5,770,214; deposited at PHLS, NCTC, 61 Colindale Avenue, London NW9 5HT under Accession No. NCTC 12165), and *S. typhi* Δ aroA/ Δ aroD (U.S. Patent No. 5,770,214;

deposited at PHLS, NCTC, 61 Colindale Avenue, London NW9 5HT under Accession No. NCTC 122309). It has been shown that one of these, CVD908-htrA, is safe and immunogenic in phase I (Tacket *et al.*, Infection and Immunity 65:452-456, 1997) and phase II studies in a total of 100 adult volunteers.

Specific examples of *S. typhimurium* mutant strains that can be used in the invention include BRD509 *S. typhimurium* Δ aroA/ Δ aroD (Strugnell *et al.*, Infection and Immunity 60:3994-4002, 1992), BRD807 *S. typhimurium* Δ aroA/ Δ htrA (Chatfield *et al.*, Microbial Pathogenesis 12:145-151, 1992; U.S. Patent No. 5,804,194; deposited at PHLS, NCTC, 61 Colindale Avenue, London NW9 5HT under Accession No. NCTC 12459), BRD698 (U.S. Patent No. 5,804,194; deposited at PHLS, NCTC, 61 Colindale Avenue, London NW9 5HT under Accession No. NCTC 12457), and BRD726 (U.S. Patent No. 5,804,194; deposited at PHLS, NCTC, 61 Colindale Avenue, London NW9 5HT under Accession No. NCTC 12458).

Additional examples of Salmonella mutant strains that can be used in the invention are described in the following publications: double *aro* mutants (WO 89/05856, U.S. Patent No. 5,770,214), *htrA* mutants (WO 91/15572, U.S. Patent No. 5,804,194), and *ompR* mutants (U.S. Patent No. 5,527,529). Also see, for example, Nakayama *et al.*, BioTechnology 6:693, 1988 and WO 92/11361. In addition, there are numerous alternative strains of *S. typhi* and *S. typhimurium* described in the literature or known in the art that are also attenuated in their virulence, and have been shown to induce immune responses against heterologous antigens. Any of these strains can be used in the method of the present invention.

Any of the attenuated *Salmonella* strains described above, or others, can be used in the method of the invention to administer a *Helicobacter* antigen to a mammal for vaccination against *Helicobacter* infection. This can be accomplished by introducing into the attenuated *Salmonella* strain a nucleic molecule encoding a *Helicobacter* antigen. The antigen-encoding nucleic acid molecule to be introduced into the attenuated *Salmonella* strain can be present, for example, in a plasmid vector (*e.g.*, pHUR3, pHUR4, pNUR3, or pNUR4 (see below)) that includes a regulatory sequence, such as a promoter, and, optionally, a sequence encoding a secretion signal (*e.g.*, a bacterial hemolysin (hly) secretion signal; WO 87/06953, U.S. Patent No. 5,143,830).

The promoter can be a prokaryotic promoter, for example, a *Salmonella* promoter, which directs expression of the *Helicobacter* antigen in the *Salmonella* vector. Examples of such promoters include the *htrA* promoter (WO 95/20665), the *nirB* promoter (WO 92/15689, U.S. Patent No. 5,547,664), the *ssaH* promoter (Valdivia *et al.*, Science 277:2007-2011, 1997), the *ompR* promoter, and any other *Salmonella* or other bacterial promoter that is upregulated when *Salmonella* is taken up by mammalian cells. Alternatively, the promoter can be a eukaryotic promoter, such as the cytomegalovirus promoter. Use of such promoters allows for expression of target antigen in a eukaryotic cell, with *Salmonella* acting as the delivery vehicle for this DNA immunization approach. The construction of such vectors is known in the art. Of course, numerous eukaryotic promoters are known in the art and can be used in the invention.

Introduction of a plasmid into an attenuated *Salmonella* strain can be accomplished using any of a number of standard methods, such as electroporation or bacteriophage transduction (Turner *et al.*, Infection and Immunity 61:5374-5380, 1993). Also see, *e.g.*, Ausubel *et al.*, *Current*

Protocols in Molecular Biology, John Wiley & Sons Inc., 1994, and Ward *et al.*, *Infection and Immunity* 67(5):2145-2152, 1999, for methods of introducing plasmids into bacteria, such as *Salmonella*.

Helicobacter Antigens

5 Preferred antigens for use in the invention are *Helicobacter* (*e.g.*, *H. pylori* or *H. felis*) proteins (*i.e.*, peptides or polypeptides), other components *Helicobacter* (*e.g.*, lipopolysaccharides, carbohydrates, or nucleic acid molecules), or immunogenic fragments thereof. Preferably, the same or a similar (*e.g.*, a fragment) antigen is used in the mucosal administration step
10 as in the parenteral administration step, however, the antigen used in each of these steps can differ. Also, preferably, the mucosally administered antigen primes an immune response to the antigen, and the parenterally administered antigen boosts an immune response to the same antigen. For the mucosal administration step, a nucleic acid molecule (*e.g.*, a DNA
15 molecule) encoding a desired antigen is inserted into an attenuated *Salmonella* vector, as is described above. For the parenteral administration step, the antigen can be, for example, purified from a bacterial culture or produced using standard recombinant or chemical synthetic methods. Methods for identifying immunogenic fragments of polypeptide antigens
20 are known in the art, and can be employed in preparing antigens for use in the method of the invention (see, *e.g.*, Sturniolo *et al.*, *Nature Biotechnology*, "Generation of Tissue-Specific and Promiscuous HLA Ligand Databases Using DNA Microarrays and Virtual HLA Class II Matrices," June, 1999). Additional antigens that can be used in the
25 parenteral administration step are whole *Helicobacter* bacteria and non-purified protein preparations, such as *Helicobacter* lysates.

The antigens used in the invention can be produced as fusion proteins, which are polypeptides containing amino acid sequences corresponding to two or more proteins (or fragments thereof) that are normally separate proteins, linked together by a peptide bond(s). Fusion proteins generally are synthesized by expression of a hybrid gene, containing nucleotides encoding each of the individual polypeptides that make up the fusion protein. An example of an antigenic fusion protein that can be used in the invention is one that contains a cholera toxin (CT) or an *E. coli* heat-labile toxin (LT) adjuvant (*e.g.*, a toxin A or B subunit, or a fragment or derivative thereof having adjuvant activity) fused to an *H. pylori* antigen, *e.g.*, a urease antigen. Another type of fusion protein included in the invention consists of an antigen fused to a polypeptide (*e.g.*, glutathione S-transferase (GST)) that facilitates purification of the fusion protein. Still another type of fusion protein that can be used in the invention is a fusion with a polypeptide that targets the expressed protein to cells of the immune system. For example, fusions with CD4 or Staph A can be used. Proteins used as antigens in the invention can also be covalently coupled or chemically cross-linked to adjuvants, using standard methods.

The most preferred *H. pylori* antigens for use in the invention are urease antigens, which include, *e.g.*, immunogenic fragments or subunits (*e.g.*, UreA or UreB) of urease. Most preferred urease antigens are enzymatically inactive, recombinant multimeric urease complexes, produced as described in Lee *et al.*, WO 96/33732. A number of other immunogenic *H. pylori* antigens can be administered according to the invention, *e.g.*, catalase (WO 95/27506), HspA and HspB (WO 94/26901), lactoferrin receptor (WO 97/13784), p76 (WO 97/12908), p32 (WO 97/12909), BabA and BabB (WO 97/47646), AlpA (WO 96/41880), AlpB (WO 97/11182), as well as the antigens described in WO 96/38475, WO

96/40893, WO 97/19098, WO 97/37044, WO 98/18323, WO 97/37044, WO 97/4764, WO 98/04702, and WO 98/32768. Additional preferred antigens for use in the invention are GHPO 1516, GHPO 789, GHPO 386, GHPO 1615, GHPO 1360, GHPO 1320, GHPO 639, GHPO 792, GHPO 536, GHPO 525, GHPO 1275, GHPO 1688, GHPO 706, GHPO 419, GHPO 1595, GHPO 1398, GHPO 986, GHPO 1282, GHPO 1056, GHPO 1443, GHPO 13, GHPO 109, GHPO 257, GHPO 1034, GHPO 236, GHPO 1166, GHPO 1351, and GHPO 1420 (WO 98/21225, WO 98/43478, and WO 98/43479), as well as other antigens described in these publications.

10 Adjuvants

Although not required, the attenuated *Salmonella* vectors described above for mucosal administration step can be administered with a mucosal adjuvant. The adjuvant can be admixed with the *Salmonella* vector or expressed in the *Salmonella* vector (*e.g.*, as a fusion protein with an antigen, see above), either from an integrated nucleic acid molecule or episomally, *e.g.*, on a plasmid. Such adjuvants can be chosen from bacterial toxins, *e.g.*, the cholera toxin (CT), the *E. coli* heat-labile toxin (LT), the *Clostridium difficile* toxin, and the Pertussis toxin (PT), or combinations, subunits, toxoids, fragments, homologs, derivatives, fusions, or mutants that are derived therefrom and have adjuvant activity. For example, it is possible to use a purified preparation of the native cholera toxin B subunit (CTB) or a polypeptide including the carboxyl-terminal repeats of *C. difficile* toxin A (WO 97/02836). Preferably, a mutant is used in which toxicity is reduced. Such mutants are described in, *e.g.*, WO 95/17211 (mutant CT Arg-7-Lys), WO 96/6627 (mutant LT Arg-192-Gly), and WO 95/34323 (mutant PT Arg-9-Lys and Glu-129-Gly). Other LT mutants that can be used include at least one of the following mutations: Ser-63-Lys, Ala-69-Gly,

Glu-110-Asp, and Glu-112-Asp. Other compounds, such as MPLA, PLGA, and QS-21, can also be used as adjuvants for the mucosal route.

Adjuvants for use in parenteral administration include, for example, aluminum compounds (*e.g.*, alum), such as aluminum hydroxide, aluminum phosphate, and aluminum hydroxy phosphate. The antigen can be precipitated with, or adsorbed onto, the aluminum compound using standard methods.

In addition to aluminum compounds, a large number of appropriate adjuvants for administration by the systemic or parenteral route exist in the art and can be used in the invention. For example, liposomes; ISCOMS; microspheres; protein choleates; vesicles consisting of nonionic surfactants; cationic amphiphilic dispersions in water; oil/water emulsions; muramidyl dipeptide (MDP) and its derivatives, such as glucosyl muramidyl dipeptide (GMDP), threonyl-MDP, murametide, and murapalmitin; QuilA and its subfractions; as well as various other compounds, such as DC-chol; monophosphoryl-lipid A (MPLA) major lipopolysaccharide from the wall of a bacterium, for example, *E. coli*, *S. minnesota*, *S. typhimurium*, *Shigella flexneri*, or *N. meningitidis*; alga-glucan; gamma-inulin; calcitriol; and loxoribine can be used. Other adjuvants, such as RIBI (ImmunoChem, Hamilton, MT) and polyphosphazene (WO 95/2415), can also be used in parenteral administration.

Useful liposomes for the purposes of the present invention can be selected, for example, from pH-sensitive liposomes, such as those formed by mixing cholesterol hemisuccinate (CHEMS) and dioleoyl phosphatidyl ethanolamine (DOPE); liposomes containing cationic lipids recognized for their fusogenic properties, such as 3-beta-(N-(N',N'-dimethylamino-ethane)carbamoyl)cholesterol (DC-chol) and its equivalents, which are

described in U.S. Patent No. 5,283,185 and WO 96/14831;
dimethyldioctadecylammonium bromide (DDAB) and the BAY compounds
described in EP 91645 and EP 206 037, for example, Bay R1005 (N-(2-
deoxy-2-L-leucylamino-beta-D-glucopyranosyl)-N-octa-
5 decyldodecanoylamide acetate; and liposomes containing MTP-PE, a
lipophilic derivative of MDP (muramidyl dipeptide). These liposomes are
useful as adjuvants with all of the antigens described herein.

Useful ISCOMs for the purposes of the present invention can be
selected, for example, from those compounds of QuilA or of QS-21
10 combined with cholesterol and, optionally, also with a phospholipid, such as
phosphatidylcholine. These are particularly advantageous for the
formulation of the lipid-containing antigens.

Useful microspheres for the purposes of the present invention can be
formed, for example, from compounds such as polylactide-co-glycolide
15 (PLAGA), alginate, chitosan, polyphosphazene, and numerous other
polymers.

Useful protein choleates for the purposes of the present invention
can be selected, for example, from those formed from cholesterol and,
optionally, an additional phospholipid, such as phosphatidylcholine. These
20 are especially advantageous for the formulation of the lipid-containing
antigens.

Useful vesicles consisting of nonionic surfactants for the purposes of
the present invention can be, for example, formed by a mixture of 1-mono-
palmitoyl glycerol, cholesterol, and dicetylphosphate. They are an
25 alternative to conventional liposomes, and can be used for the formulation
of all of the antigens described herein.

Useful oil/water emulsions for the purposes of the present invention can be selected, for example, from MF59 (Biocine-Chiron), SAF1 (Syntex), and the montanides ISA51 and ISA720 (Seppic).

A useful adjuvant for the purposes of the present invention can also
5 be a fraction derived from the bark of the South American tree *Quillaja Saponaria Molina*, for example, QS-21, a fraction purified by HPLC chromatography as is described in U.S. Patent No. 5,057,540. Since some toxicity may be associated with QS-21, it may be advantageous to use it in liposomes based on sterol, as is described in WO 96/33739.

10 Induction of an Immune Response Against Helicobacter

The method of the invention can be used to prevent *Helicobacter* infection in a patient, as well as to treat an ongoing *Helicobacter* infection in a patient. Thus, gastroduodenal diseases associated with these infections, including acute, chronic, or atrophic gastritis, and peptic ulcers, *e.g.*, gastric
15 or duodenal ulcers, can be prevented or treated using the method of the invention.

As is noted above, the method of the invention involves mucosal (*e.g.*, oral, intranasal, intragastric, pulmonary, intestinal, rectal, ocular, vaginal, or urinary tract) administration of a *Salmonella* vector including a
20 nucleic acid molecule that encodes a *Helicobacter* antigen, followed by parenteral (*e.g.*, intramuscular, subcutaneous, intradermal, intraepidermal, intravenous, or intraperitoneal) administration of a *Helicobacter* antigen, preferably in association with an adjuvant. The antigen used in the mucosal prime can be different from, cross-reactive with, or, preferably, identical to
25 the parenterally administered antigen. Preferably, the mucosal administration step primes an immune response to an antigen, and the parenteral administration step boosts an immune response to the antigen.

Also included in the invention are vaccination methods involving parenteral priming and mucosal boosting (*e.g.*, with a *Salmonella* vector including a nucleic acid molecule encoding a *Helicobacter* antigen), and parenteral administration of a *Salmonella* vector including a nucleic acid molecule encoding a *Helicobacter* antigen.

Attenuated *Salmonella* vectors, antigens, formulations, adjuvants, administration regimens, specific mucosal and parenteral routes, and dosages to be used in the method of the invention can readily be determined by one skilled in the art. For example, 5×10^6 - 5×10^{10} colony forming units, *e.g.*, 5×10^8 colony forming units, of an attenuated *Salmonella* vector can be used in the mucosal administration, and 5-1000 μg , *e.g.*, 100 μg , antigen, can be used in the parenteral administration. The mucosal administration can take place only once or two or more (*e.g.*, three, four, or five) times, for example, separated by two, three, or four days or weeks. Similarly, the parenteral administration can take place once or two or more (*e.g.*, three, four, or five) times, separated by weeks, months, or years from each other or the mucosal administration.

In one example of an immunization regimen that can be used, a patient is primed with two doses of an attenuated *Salmonella* vector (*e.g.*, *S. typhi* CVD908-htrA or CVD908, or *S. typhimurium* BRD509 or BRD807) expressing an antigen (*e.g.*, urease from plasmid pHUR3, pHUR4, pNUR3, or pNUR4) on days 0 and 21, and then parenterally boosted on day 51 or later with an antigen (*e.g.*, urease) and an adjuvant (*e.g.*, alum). The details of construction of pHUR3 and pNUR3, which each include an ampicillin resistance gene, are described below. pHUR4 and pNUR4 are constructed by removing the ampicillin resistance gene from pHUR3 and pNUR3, respectively, by digestion with the restriction endonuclease *RcaI*, and

cloning into the digested vectors a kanamycin resistance gene that can be obtained from plasmid pUC4K (Pharmacia) by digestion with *EcoRI*.

A useful pharmaceutical composition for the purposes of the present invention can be manufactured in a conventional manner. In particular, it
5 can be formulated with a pharmaceutically acceptable carrier or diluent, *e.g.*, water or a saline solution. In general, the diluent or carrier can be selected according to the mode and route of administration and according to standard pharmaceutical practices. Appropriate carriers or diluents, as well
10 as what is essential for the preparation of a pharmaceutical composition, are described, *e.g.*, in *Remington's Pharmaceutical Sciences* (18th edition), ed. A. Gennaro, 1990, Mack Publishing Company, Easton, PA., a standard reference book in this field. As a specific example, the attenuated *Salmonella* vectors of the invention can be formulated in a tablet for oral
administration (see, *e.g.*, U.S. Patent No. 5,804,194).

15 The therapeutic or prophylactic efficacy of the method of the invention can be evaluated according to standard methods, *e.g.*, by measuring the induction of an immune response or the induction of therapeutic or protective immunity using, *e.g.*, the mouse/*H. felis* model and the procedures described in Lee *et al.*, Eur. J. Gastroenterology and
20 Hepatology 7:303, 1995 or Lee *et al.*, J. Infect. Dis. 172:161, 1995. Persons skilled in this art will realize that *H. felis* can be replaced in the mouse model by another *Helicobacter* species. For example, the efficacy of the method is, preferably, evaluated in a mouse model using an *H. pylori* strain adapted to mice. The efficacy can be determined by comparing the level of
25 infection in gastric tissue (*e.g.*, by measuring the urease activity, bacterial load, or condition of the gastritis) with that in a control group. A therapeutic effect or a protective effect exists when infection is reduced

compared with a control group. Experimental methods and results showing the efficacy of the present method is described as follows.

Experimental methods and results

Construction of ureAB expression plasmids under the control of the nirB and htrA promoters - Method 1

A *ureAB* expression plasmid is constructed by subcloning a PCR product containing the *ureAB* genes (amplified from plasmid pORV273) into plasmid vector ptetnir15. Plasmid pORV273 is obtained from OraVax, Inc., Cambridge, MA. Plasmid ptetnir15 has been described (Chatfield *et al.*, Bio/Technology 10:888-892, 1992; Oxeer *et al.*, Nucl. Acids Res. 19:1889-1892, 1991). This vector was modified by standard techniques known in the art, to introduce into the vector a suitable restriction site for subcloning other genes for optimal expression under control of the *nirB* promoter. An *NcoI* site was introduced 10 basepairs 3' to the Shine-Dalgarno sequence of ptetnir15, and the resultant plasmid is designated ptetnir15/mod. Plasmid ptetnir15/mod, carried in strain BRD940, is obtained from Peptide Therapeutics Ltd., Cambridge, U.K.

The *ureAB* gene is amplified by PCR from pORV273 using Turbo Pfu polymerase (Stratagene), which has 3'-5' proof-reading activity, and two primers, designated orafor and orarev. Primer orafor introduces *EcoRI* and *BspHI* sites immediately upstream of the initiating codon of the *ureA* gene. Primer orarev binds approximately 18 basepairs downstream of the *BamHI* site that is located 45 basepairs downstream of the termination codon of the *ureB* gene.

The PCR reaction includes 0.1 µg pORV273 and 100 pmol each of primers orafor (5'-TAG GGA ATT CTC ATG AAA CTC ACC CCA AAA G-3' (SEQ ID NO:31)) and orarev (5'-GCC AAC TTA GCT TCC TTT

CGG G-3' (SEQ ID NO:32)) per 100 µl reaction and utilizes 25 cycles, with an annealing temperature of 50°C. The resulting 2.4 kb PCR product is purified from a 1% agarose gel using a Qiaquick gel extraction kit (Qiagen).

As is described below, the actual method used in the generation of pNUR and pHUR differed from this description in the sequence of orarev.

Therefore, the method described here may need to be adapted in ways known to those skilled in the art by changing, for example, the precise annealing temperature or the number of cycles required to give sufficient product, or even in the sequence of the primer orarev.

The PCR product is digested with *Bsp*HI + *Bam*HI, and purified with a Promega Wizard DNA clean-up kit. Plasmid ptetnir15/mod is digested with *Nco*I + *Bam*HI (the *Nco*I site is 10 basepairs 3' to the Shine-Dalgarno sequence of ptetnir15, and generates a cohesive end that is compatible with *Bsp*HI), and dephosphorylated using shrimp alkaline phosphatase. The largest fragment from the digestion of ptetnir15/mod is isolated from a 1% agarose gel using a Qiaquick gel extraction kit (Qiagen), and ligated to the digested PCR product using the Ligator Express Kit (Clontech). Ligations are transformed into electrocompetent *E. coli* TG1 cells (Stratagene).

Plasmids from ampicillin-resistant transformants are screened for the presence of the full length, 2.4 kb *ureAB* gene by restriction analysis. The *ureAB* gene from plasmid pORV273 has a *Bam*HI site within the coding sequence. However, in a small number of ptetnir15/mod + *ureAB* transformants, incomplete digestion or re-ligation of the two *ureAB* fragments yields the full length *ureAB* PCR product. The orientation of the *ureAB* gene in the ptetnir15-derived plasmid is confirmed by PCR, and a plasmid with the full length *ureAB* gene, in the correct orientation is designated pNUR.

The *nirB* promoter in plasmid pNUR is replaced with the *htrA* promoter from phtrAcore, which is obtained from Peptide Therapeutics Ltd., Cambridge, U.K. Plasmids pNUR and phtrAcore are digested with *Pst*I and *Bgl*II. Digested pNUR is dephosphorylated with shrimp alkaline phosphatase. The digestion products are run on a 1% agarose gel, and a 0.8 kb fragment containing the *htrA* promoter from the phtrAcore digestion and the 4.0 kb fragment from pNUR lacking the *nirB* promoter are extracted from the gel using a Qiagen Qiaquick gel extraction kit. The two fragments are ligated together (Clontech Ligator express kit), and transformed into electrocompetent *E. coli* TG1 cells (Stratagene). Transformants are screened for the presence of the *htrA* promoter by PCR using primer pairs specific for *htrA* (5902/5904) or *nirB* (5901/5904). A plasmid with the *htrA* promoter and a full length *ureAB* gene is designated pHUR.

The nucleotide sequence across the promoter region and *ureAB* genes of final plasmids are confirmed. Samples of the plasmids are prepared using the Qiagen "Plasmid midi kit" (Catalog No. 12143), and the DNA sequence determined by standard techniques. Oligonucleotides 5901 to 5919 (see below) can be used, and allow nucleotide sequence determination of both DNA strands. Oligonucleotides 5901 and 5902 hybridize within *nirB* and *htrA*, respectively, while 5919 hybridizes within ptdnir15/mod, downstream of the *ureAB* genes. The other oligonucleotides hybridize within the *ureAB* genes. The data confirm that the nucleotide sequence across the recombinant region of all plasmids are as expected.

Plasmids pNUR and pHUR are introduced into *S. typhimurium* strains such as, e.g., BRD509 and BRD807, and *S. typhi* strains such as, e.g., CVD908 and BRD948, by electroporation and selection of ampicillin-resistant colonies.

Construction of ureAB expression plasmids under the control of the nirB and htrA promoters - Method 2

The protocol described above is one example of many by which one skilled in the art can derive an expression plasmid suitable for directing the synthesis of an *H. pylori* antigen, *e.g.*, urease, under the control of the *htrA* or *nirB* promoter in an attenuated strain of *Salmonella*. Alternative primers can be used in the PCR amplification of the genes from the starting plasmid, and alternative strategies for the introduction of the gene *via* alternative restriction sites are possible. One such alternative was employed in the construction of plasmids pNUR3 and pHUR3. During the design of the primers for PCR, a sequence error in the database-deposited gene sequence caused the 3' end of the *ureB* gene to be incorrectly identified. A primer was synthesized for the PCR amplification that, in fact, resulted in a non-native sequence of the gene, containing an additional 49 codons after the genuine termination codon. This error was subsequently corrected by the method described below, yielding a final plasmid with a sequence identical to that of the plasmid that would be produced by the strategy described above. This method is described in further detail, as follows.

As is described above, plasmid pORV273 was obtained from OraVax Inc. Plasmid ptnir15 has been described (Chatfield *et al.*, Bio/Technology 10:888-892, 1992; Oxeer *et al.*, Nucl. Acids Res. 19:1889-1892, 1991), and this vector was modified by standard techniques, to introduce into the vector a suitable restriction site for subcloning other genes for optimal expression under control of the *nirB* promoter. An *NcoI* site was introduced 10 basepairs 3' to the Shine-Dalgarno sequence of ptnir15, and the resultant plasmid was designated ptnir15/mod. Plasmid ptnir15/mod, carried in strain BRD940, was obtained from the culture collection of Peptide Therapeutics Ltd., Cambridge, U.K.

The *ureAB* gene was amplified by PCR from pORV273 using Turbo Pfu polymerase (Stratagene), which has 3'-5' proof-reading activity and two primers, designated orafor and orarev. Primer orafor introduces *EcoRI* and *BspHI* sites immediately upstream of the initiating codon of the *ureA* gene. Primer orarev introduces a *BamHI* and a *PstI* site just before the correct 3' end of the *ureAB* gene. Subsequent digestion and cloning, as is described below, resulted in the deletion of the correct termination codon of *ureB*, with the result that transcription continued into the vector sequence until an in-frame stop codon was reached, adding 49 amino acids to the translated protein.

The PCR reaction included 0.1 µg pORV273 and 100 pmol each of primers orafor (5'-TAG GGA ATT CTC ATG AAA CTC ACC CCA AAA G-3' (SEQ ID NO:31)) and orarev (5'-TCT ACT GCA GGA TCC AAA ATG CTA AAG AGT TGC G-3' (SEQ ID NO:33)) per 100 µl reaction, and utilized 25 cycles, with an annealing temperature of 50°C. The resulting 2.4 kb PCR product was purified from a 1% agarose gel using a Qiaquick gel extraction kit (Qiagen). The PCR product was digested with *BspHI* + *BamHI*, and purified with a Promega Wizard DNA clean-up kit. Plasmid ptnir15/mod was digested with *NcoI* + *BamHI* (the *NcoI* site is 10 basepairs 3' to the Shine-Dalgarno sequence of ptnir15, and generates a cohesive end that is compatible with *BspHI*), and dephosphorylated using shrimp alkaline phosphatase. The largest fragment from the digestion of ptnir15/mod was isolated from a 1% agarose gel using a Qiaquick gel extraction kit (Qiagen), and ligated to the digested PCR product using the Ligator Express Kit (Clontech). Ligations were transformed into electrocompetent *E. coli* TG1 cells (Stratagene).

Plasmids from ampicillin-resistant transformants were screened for the presence of the full length, 2.4 kb *ureAB* gene by restriction analysis. The *ureAB* gene from plasmid pORV273 has a *Bam*HI site within the coding sequence. However, in a small number of ptnir15/mod + *ureAB* transformants, incomplete digestion or re-ligation of the two *ureAB* fragments yielded the full length *ureAB* PCR product. The orientation of the *ureAB* gene in the ptnir15-derived plasmid was confirmed by PCR and a plasmid with the full length *ureAB* gene, in the correct orientation was designated pNUR1.

The *nirB* promoter in plasmid pNUR1 was replaced with the *htrA* promoter from phtrAcore, which is obtained from Peptide Therapeutics Ltd., Cambridge, U.K. Plasmids pNUR1 and phtrAcore were digested with *Pst*II and *Bgl*II. Digested pNUR1 was dephosphorylated with shrimp alkaline phosphatase. The digests were run on a 1% agarose gel, and a 0.8 kb fragment containing the *htrA* promoter from the phtrAcore digest and the 4.0 kb fragment from pNUR1 lacking the *nirB* promoter were extracted from the gel using a Qiagen Qiaquick gel extraction kit. The two fragments were ligated together (Clontech Ligator express kit) and transformed into electrocompetent *E. coli* TG1 cells (Stratagene). Transformants were screened for the presence of the *htrA* promoter by PCR using primer pairs specific for *htrA* (5902/5904) or *nirB* (5901/5904). A plasmid with the *htrA* promoter and a full length *ureAB* gene was designated pHUR1.

Subsequent to this it was discovered that there had been a cloning error in the 3' terminal portion of *ureB*, resulting in a translated product with an additional 49 amino acids from both pHUR1 and pNUR1. This was corrected by replacing the small *Bam*HI fragment containing the C-terminus of the *ureB* gene with the corresponding, and correct, fragment from pORV272. pORV273, pHUR1, and pNUR1 were digested with *Bam*HI,

and the small fragment from the pORV273 digestion was ligated to the large fragment from the pHUR1 and pNUR1 digestions. Clones were screened for orientation of the insert, and clones with the correct orientation were designated pHUR3 and pNUR3. These clones were characterized by full nucleotide sequencing of the region including the promoter and the complete *ureAB* gene on both strands, and found to be correct.

The nucleotide sequences across the *nirB* promoter and *ureAB* genes of pNUR1 and of the *htrA* promoter region of pHUR1 were confirmed. Samples of the two plasmids were prepared using the Qiagen "Plasmid midi kit" (Catalogue No. 12143), and the DNA sequence was determined by standard techniques known in the art. Oligonucleotides 5901 to 5919 were used, which allow nucleotide sequence determination of both DNA strands. Oligonucleotides 5901 and 5902 hybridize within *nirB* and *htrA*, respectively, while 5919 hybridizes within *ptetnir15/mod* downstream of the *ureAB* genes. The other oligonucleotides hybridize within the *ureAB* genes. These were diluted to 1 pmol μl^{-1} , packed in dry ice with the plasmid samples, and sent to Cambridge Bioscience (Cambridge) for nucleotide sequence determination. The data confirmed that the nucleotide sequence across the recombinant region of all three plasmids was as expected.

Sequences of primers that can be used in the invention, as is described above, are as follows.

5901

Primes within *nirB* promoter ~60 basepairs upstream of SD sequence
TCA AAT GGT ACC CCT TGC TGA (SEQ ID NO:34)

5902

Primes within *htrA* promoter ~60 basepairs upstream of SD sequence
TAT TCC GGA ACT TCG CGT TA (SEQ ID NO:35)

5903

Primes ~250 basepairs downstream from start of *ureA* gene

TGT TTC CTG ATG GGA CTA AAC TC (SEQ ID NO:36)

5904

5 Reverse primes ~300 basepairs downstream from start of *ureA* gene

ACC AGG AAC TAA TTT ACC ATT G (SEQ ID NO:37)

5905

Primes ~550 basepairs downstream from start of *ureA* gene

TTG ATT GAC ATT GGC GGT AAC (SEQ ID NO:38)

10

5906

Reverse primes ~600 basepairs from start of *ureA* gene

GTT GTC TGC TTG TCT ATC AAC C (SEQ ID NO:39)

5907

Primes ~150 basepairs downstream from start of *ureB* gene

15 GGT GGC GGT AAA ACC CTA AGA G (SEQ ID NO:40)

5908

Reverse primes ~180 basepairs downstream of *ureB* gene

CTT TGC TAG GGT TGT TAG ATT G (SEQ ID NO:41)

5909

20 Primes ~400 basepairs downstream from start of *ureB* gene

AAT CCC TAC AGC TTT TGC AAG C (SEQ ID NO:42)

5910

Reverse primes ~500 basepairs from start of *ureB* gene

GTG CCA TCA GCA GGA CCG GTT C (SEQ ID NO:43)

25

5911

Primes ~750 basepairs from start of *ureB* gene

ATC GCC ACA GAC ACT TTG AAT G (SEQ ID NO:44)

5912

Reverse primes ~820 basepairs downstream from start of *ureB* gene

30 TAG CAG CCA TAG TGT CTT CTA C (SEQ ID NO:45)

- 5913
Primes ~1050 basepairs downstream from start of *ureB* gene
TGA AGA CAC TTT GCA TGA CAT G (SEQ ID NO:46)
- 5914
5 Reverse primes 1080 basepairs downstream of *ureB* gene
TGA GAG TCA GAA CTG GTG ATT G (SEQ ID NO:47)
- 5915
Primes ~1350 basepairs downstream from start of *ureB* gene
CAT GAT CAT CAA AGG CGG ATT C (SEQ ID NO:48)
- 10 5916
Reverse primes ~1380 basepairs downstream from start of *ureB*
GAA GCG TTC GCA TCG CCC ATT TG (SEQ ID NO:49)
- 5917
Primes ~1650 basepairs from start of *ureB*
15 TCG TGG ATG GCA AAG AAG TAA C (SEQ ID NO:50)
- 5918
Reverse primes ~1680 basepairs from start of *ureB*
GCG CCA AGC TCA CTT TAT TG (SEQ ID NO:51)
- 5919
20 Reverse primes ~80 basepairs downstream of *Bam*HI site downstream of
ureB
CAA CGA CAG GAG CAC GAT CAT G (SEQ ID NO:52)

25 The nucleotide sequences across the promoter regions and *ureAB* genes of the final plasmids, pHUR3 and pNUR3, were also confirmed. *E. coli* MC1061 cells containing the plasmids were sent to Cambridge Biosciences Ltd., who prepared plasmid DNA and determined the nucleotide sequences of the promoter and *ureAB* genes of both plasmids. The data confirmed that the nucleotide sequence across the relevant region of both plasmids was as expected. The sequence of plasmid pHUR3 is

shown in Fig. 4, and a plasmid map showing its relevant features is provided in Fig. 5.

Plasmids pNUR and pHUR were introduced into *S. typhimurium* strains BRD509 and BRD807, and *S. typhi* strains CVD908 and BRD948, by electroporation and selection of ampicillin-resistant colonies.

Immunization and Protection Experiments

Inbred Balb/C mice were immunized by the intragastric route with live, attenuated *Salmonella typhimurium* (1E10 CFU/ml) expressing urease apoenzyme on day 0 (Fig. 1). Animals were boosted twice on days 21 and 35 with 10 µg soluble, recombinant urease plus aluminum hydroxide (200 µg) by the parenteral route. Fourteen days later, serum antibody responses to urease were measured. Controls included: (1) prime-boost with the *Salmonella* parental control strains (BRD509 Δ aroA/ Δ aroD (Strugnell *et al.*, Infection and Immunity 60:3994-4002, 1992) and BRD807 Δ aroA/ Δ htrA (Chatfield *et al.*, Microbial Pathogenesis 12:145-151, 1992)) minus the urease construct, (2) mucosal priming with LT in place of *Salmonella* (gold standard), and (3) parenteral immunization with urease plus alum alone. Attenuated *S. typhimurium* (Δ aroA/ Δ aroD) expressing urease under the transcriptional control of either an *htrA* promoter (pHUR3) or the *nirB* promoter (pNUR3) induced an elevated IgG2a response against urease that was greater than the gold standard using LT-Alum (Fig. 2A). A comparable response to LT-Alum was induced with *S. typhimurium* (Δ aroA/ Δ htrA) carrying the same urease constructs (Fig. 2A). Analysis of the IgG1/IgG2a ratio demonstrated the induction of a Th1 response with the double aro mutant, and a more balanced response with the Δ aro/ Δ htrA mutant strain (Fig. 2B). Urease-specific antibody in Fig. 2A is expressed as EU/ml on a logarithmic scale and median response is indicated by the bar.

The level of protective efficacy employing *S. typhimurium*-vectored urease in a prime-boost strategy was determined. Fig. 3A shows the results of quantitative *H. pylori* culture of mice immunized on day 0 with 1E10 CFU/ml live attenuated *S. typhimurium* (Δ aroA/ Δ aroD or Δ aroA/ Δ htrA) and boosted on days 21 and 35 with urease (10 μ g) plus alum (200 μ g). Three weeks later, animals were challenged with *H. pylori* (1E7 CFU/ml) and efficacy was assessed in gastric tissue 4 weeks later using quantitative culture. Strains including the urease constructs are indicated in the key of Fig. 3A. Fig. 3B shows protection depicted as log₁₀ reduction in comparison to the no treatment (Tx) control group. A significant reduction in bacterial burden was observed when attenuated Salmonella expressing urease was administered as part of a prime-boost regimen with alum (Wilcoxon rank sum compared to parental control strain). No significant difference was observed between group 1 (pHUR3-Alum) and group 7 (LT-Alum).

All patents and publications cited above are hereby incorporated by reference in their entirety.

What is claimed is:

1. A method of inducing an immune response against *Helicobacter* in a mammal, said method comprising the steps of:

mucosally administering to said mammal an attenuated *Salmonella* vector comprising a nucleic acid molecule encoding a *Helicobacter* antigen,
5 and

parenterally administering to said mammal a *Helicobacter* antigen.

2. The method of claim 1, wherein said attenuated *Salmonella* vector is administered orally to said mammal.

3. The method of claim 1, wherein said *Helicobacter* antigen is a
10 urease, a urease subunit, or an immunogenic fragment thereof

4. The method of claim 1, wherein said mammal is at risk of developing, but does not have, a *Helicobacter* infection.

5. The method of claim 1, wherein said mammal has a *Helicobacter* infection.

15 6. The method of claim 1, wherein said parenteral administration of said *Helicobacter* antigen further includes parenteral administration of an adjuvant.

7. The method of claim 6, wherein said adjuvant is an aluminum compound.

20 8. The method of claim 7, wherein said aluminum compound is alum.

9. The method of claim 1, wherein said attenuated *Salmonella* vector is a *Salmonella typhi* vector.

10. The method of claim 9, wherein said *Salmonella typhi* vector is CVD908-htrA or CVD908.

5 11. The method of claim 1, wherein the attenuated *Salmonella* vector is a *Salmonella typhimurium* vector.

12. The method of claim 11, wherein said *Salmonella typhimurium* vector is BRD509 or BRD807.

10 13. The method of claim 1, wherein said attenuated *Salmonella* vector further comprises an *htrA* promoter.

14. The method of claim 1, wherein said attenuated *Salmonella* vector further comprises a *nirB* promoter.

15 15. The method of claim 1, wherein said mucosal administration primes an immune response to an antigen and said parenteral administration boosts an immune response to said antigen.

16. An attenuated *Salmonella* vector comprising a nucleic acid molecule encoding a *Helicobacter* antigen.

17. The vector of claim 16, wherein said antigen is a urease, a urease subunit, or an immunogenic fragment thereof.

18. The vector of claim 16, wherein said attenuated *Salmonella* vector is a *Salmonella typhi* vector.

19. The vector of claim 18, wherein said *Salmonella typhi* vector is CVD908-htrA or CVD908.

5 20. The vector of claim 16, wherein the attenuated *Salmonella* vector is a *Salmonella typhimurium* vector.

21. The vector of claim 20, wherein said *Salmonella typhimurium* vector is BRD509 or BRD807.

10 22. The vector of claim 16, wherein said attenuated *Salmonella* vector further comprises an *htrA* promoter.

23. The vector of claim 16, wherein said attenuated *Salmonella* vector further comprises a *nirB* promoter.




Fig. 1

Figure 2. Mucosal priming with *S. typhimurium* vectored urease followed by parenteral boosting with alum induces an IgG2a humoral immune response equivalent to that induced by mucosal priming with LT.




Figure 3. Mucosal priming with attenuated *S. typhimurium*::*ureA/B* affords equivalent protection as LT in a prime-boost regimen with Alum




Fig. 4 (1 of 4)




Fig. 4 (2 of 4)

TGTAACAACCATGATTGGTGGTGGGAACCGGTCTGCTGATGGCACTAATGCGACTACTATCACTCCAGGCAGAAGAAATTTAAATGGAT 1350
UreB
V T T M I G G G T G P A D G T N A T T I T P G R R N L K W M
GCTCAGAGCGGCTGAAGAATATTCTATGAATTAGGTTTCTTGGCTAAAGGTAACGCTTCTAACGATCCGAGCTTAGCCGATCAAAATGA 1440
UreB
L R A A E E Y S M N L G F L A K G N A S N D A S L A D D I E
AGCCGGTGGCATTGGCTTTGCAATTCACGAAGACTGGGSCACCACTCCTTCTGCAATCAATCATGCGTTAGATGTTGCGGACAAATACGA 1530
UreB
A G A I G F A I H E D W G T T P S A I N H A L D V A D K Y D
TGTGCAAGTCGCTATGCCACAGACACTTTGAATGAAGCGGTTGTGTAGAAGACACTATGGCTGCTATTGCTGGACGCACTATGCACAC 1620
UreB
V G V A I A T D T L N E A G C V E D T M A A I A G R T M H T
TTTCCACACTGAAGGCGCTGGCGGCGACACGCTCTGATATTATTAAGTAGCGGTGAACACAACATTCTTCCGCTTCCACTAACCC 1710
UreB
F H T E G A G G G H A P D I I K V A G E H N I L P A S T N P
CACCATCCCTTTCACCGTGAATACAGAAGCAGAGCACATGGACATGCTTATGGTGTGCCACCACTTGGATAAAAGCATTAAAGAAGATGT 1800
UreB
T I P F T V N T E A E H M D M L M V C H H L D K S I K E D V
Bam HI
TCAGTTGCTGATTCAAGGATCCGCCCTCAAACCAATTGCGGCTGAAGACACTTTGCAATGACATGGGGATTTTCTCAATCACCAGTTCTGA 1890
UreB
O F A D S R I R P O T I A A E D T L H D M G I F S I T S S D
CTCTCAAGCGATGGGCGTGTGGGTGAAGTTATCACTAGAACCTTGGCAACAGCTGACAAAAACAAGAAAGAAATTTGGCCGCTTGAAGA 1980
UreB
S O A M G R V G E V I T R T W O T A D K N K K E F G R L K E
AGAAAAAGGCGATAACGACAACTTCAGGATCAACGCTACTTGTCTAAATACACCAATTAACCCAGCGATCGCTCATGGGATTAGCGAGTA 2070
UreB
E K G D N D N F R I K R Y L S K Y T I N P A I A H G I S E Y
TGTAGGTTCACTAGAAAGTGGGCAAGTGGCTGACTTGGTATTGTGGAGTCCAGCATTCCTTGGCGTGAACCAACATGATCATCAAGG 2160
UreB
V G S V E V G K V A D L V L W S P A F F G V K P N M I I K G
CGGATTCAATTGCGTTAAGCCAAATGGGCGATGCGAAGCTTCTATCCCTACCCACAACCGGTTTATTACAGAGAAATGTTGCTCATCA 2250
UreB
G F I A L S O M G D A N A S I P T P O P V Y Y R E M F A H H
TGGTAAAGCTAAATACGATGCAAAACATCACTTTTGTGCTCAAGCGGCTTATGACAAAGGCATTAAAGAAGAATTAGGACTTGAAAGACA 2340
UreB
G K A K Y D A N I T F V S O A A Y D K G I K E E L G L E R O
AGTGTGCGGTAAGAAATTCAGAAATATCACTAAAAAGACATGAATTCAACGACACTACCGCTCACATTGAAGTCAATCCTGAAAC 2430
UreB
V L P V K N C R N I T K K D M O F N D T T A H I E V N P E T
TTACCATGTCTTCGTGGATGGCAAGAAGTAACCTTAAACCAGCCAAATAAAGTGAGCTTGGCGCAACTCTTTAGCATTTTCTAGGATTT 2520
UreB
Y H V F V D G K E V T S K P A N K V S L A O L F S I F D F
Bam HI
TTTAGGAGCAACGCTCCTTAGATCCCCGGGAATTGGGGATCCGCTAGCCCGCTAATGAGCGGGCTTTTTTCTCGGGCAGCGTTGGGT 2610
L G A T L L R S P G I G D P L A R L M S G L F F L G O R W V

Fig. 4 (3 of 4)

CCTGGCCACGGGTGCGCATGATCGTCTCTGCTGTTGAGGACCCGGCTAGGCTGGCGGGTTCCTTACTGGTTAGCAGAAATGAATCAC 2700
L A T G A H D R A P V Y E D P A R L A G L P Y W L A E . I T
CGATACGCGAGCGAACGTGAAGCGACTGCTGCTGCAAAACGCTGCGACCTGAGCAACAACATGAATGGTCTTCGGTTTCCGTTTCGT 2790
D T R A N V K R L L L O N V C D L S N N M N G L R F P C F V
AAAGTCTGGAACGCGGAAGTCAGCGCTCTTCCGTTCTCTGCTCACTGCTGCTGCGTTCGGTTCGGTTCGGCGAGCGGTATCA 2880
K S G N A E V S A L P L P R S L T R C A R S F G C G E R Y O
GCTCACTCAAGGCGGTAAACGGTTATCCACAGAATCAGGGGATAACGAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGG 2970
Origin
L T O R R . Y G Y P O N O G I T O E R T C E O K A S K R P G
AACCCTAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGG 3060
Origin
T V K R P R C W R F S I G S A P L T S I T K I D A O V R G G
CGAAACCCGACAGACTATAAGATACAGGCGTTTTCCCTGGAAGCTCCCTGCTGCGCTCTCTGTTCCGACCTGCGCGTTACCGGA 3150
Origin
E T R O D Y K D T R R F P L E A P S C A L L F R P C R L P D
TACCTGTCGCGCTTCTCCCTTGGGAAGCGTGGCGCTTCTCAATGCTCAGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCTCC 3240
Origin
T C P P F S L R E A W R F L N A H A V G I S V R C R S F A P
AAGTGGGCTGTGTGCAAGACCCCGCTTACGCCGACCGCTGCGCTTATCCGGTAACATATCGTCTTGAAGTCAACCCGGTAAGACAC 3330
Origin
S W A V C T N P P F S P T A A P Y P V T I V L S P T R . D T
GACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTCTTGAAGTGGTGGCT 3420
Origin
T Y R H W O O P L V T G L A E R G M . A V L O S S . S G G L
AACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCC 3510
Origin
T T A T L E G O Y L V S A L C . S O L P S E K E L V A L D P
GGCAAAACAAACCACCGCTGCTGAGCGGTGTTTTTTGTTTGAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAACATCCTTTG 3600
Origin
A N K P P L V A V V F L F A S S R L R A E K K D L K K I L .
ATCTTTTCTACGGGCTGACGCTCAGTGGAAACGAAACCTACGTTAAGGGATTTTGGTCATGAGATTATCAAAAGGATCTTACCTAG 3690
S F L R G L T L S G T K T H V K G F W S . D Y O K G S S P R
ATCCTTTTAAATTAATAATGAAGTTTAAATCAATCTAAAGTATATAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAG 3780
AmpR
S F . I K N E V L N D S K V Y H S K L G L T V T N A . S V R
GCACTATCTCAGCGATCTGTCTATTTCTGTTATCCATAGTTGCTGACTCCCGTCTGTAGATAACTACGATACGGGAGGGCTTACCA 3870
AmpR
H L S O R S V Y F V H P . L P D S P S C R . L R Y G R A Y H
TCTGGCCCCAGTGTGCAATGATACGCGAGACCCACGCTACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAG 3960
AmpR
L A P V L O . Y R E T H A H R L O I Y D O . T S O P E G P S
CGCAGAAGTGGTCTGCACTTTATCCGCTCCATCCAGTCTATTAAATGTTGCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGT 4050
AmpR
A E V V L O L Y P P P S S L L I V A G K L E . V V R O L I V

Fig. 4 (4 of 4)

Pst I

TTGCGCAACGTTGTTGCCATTGCTGCAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCTTCATTGAGCTCCGGTTCCCAACGATCA 4140

AmpR

C A T L L P L L D A S W C H A R R L V W L H S A P V P N D O

AGGCGAGTTACATGATCCCCATGTTGTGCAAAAAGCGGTAGCTCCTTCGGTCCGATCGTTGTCAGAAGTAAGTTGCCCGCAGTG 4230

AmpR

G E L H D P P C C A K K R L A P S V L R S L S E V S W P O C

TTATCACTCATGGTTATGGCAGCACTGCATAATTCTTACTGTGATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACC 4320

AmpR

Y H S W L W O H C I I L L L S C H P . D A F L . L V S T O P

AAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAACACGGGATAATACGCGCCACATAGCAGAACTTTA 4410

AmpR

S H S E N S V C G D R V A L A R R O H G I I P R H I A E L .

AAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGT 4500

AmpR

K C S S L E N V L R G E N S O G S Y R C . D P V R C N P L V

GCACCAACTGATCTTCAGCATCTTTTACTTTCAACGAGCTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATA 4590

AmpR

H P T D L O H L L L S P A F L G E O K O E G K H P O K R E .

AGGGCGACACGGAATGTTGAATACTCATACTCTTCCTTTTCAATATTATTGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATAC 4680

AmpR

G R H G N V E Y S Y S S F F N I I E A F I R V I V S . A D T

ATATTTGAATGATTTAGAAAAATAACAAATAGGGGTTCCGCGCACATTTCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTTAT 4770

Y L N V F R K I N K . G F R A H F P E K C H L T S K K P L L

ATCATGACATTAACCTATAAAAATAGGCGTATCAGGAGCCCTTTCGTCTTCAA 4824

S . H . P I K I G V S R G P F V F K




Fig. 5

SEQUENCE LISTING

<110> OraVax, Inc.

<120> Use of salmonella vectors for
vaccination against helicobacter infection

<130> 06132/060W01

<150> US 09/431,705

<151> 1999-11-01

<160> 52

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 4824

<212> DNA

<213> Artificial Sequence

<220>

<223> includes sequences from Helicobacter pylori,
Salmonella typhimurium, and Escherichia coli

<221> CDS

<222> (2)...(31)

<221> CDS

<222> (41)...(61)

<221> CDS

<222> (65)...(799)

<221> CDS

<222> (803)...(2512)

<221> CDS

<222> (2516)...(2692)

<221> CDS

<222> (2696)...(2896)

<221> CDS

<222> (2900)...(3322)

<221> CDS

<222> (3326)...(3385)

<221> CDS
 <222> (3389)...(3406)

<221> CDS
 <222> (3410)...(3466)

<221> CDS
 <222> (3470)...(3598)

<221> CDS
 <222> (3602)...(3661)

<221> CDS
 <222> (3665)...(3697)

<221> CDS
 <222> (3701)...(3769)

<221> CDS
 <222> (3773)...(3817)

<221> CDS
 <222> (3821)...(3844)

<221> CDS
 <222> (3848)...(3889)

<400> 1
 g aat tct att ccg gaa ctt cgc gtt ata aaa tgaatctga cgt aca cag 49
 Asn Ser Ile Pro Glu Leu Arg Val Ile Lys Arg Thr Gln
 1 5 10

caa ttt aga tat taa tca tcc aca gga gag atc tcc atg aaa ctc acc 97
 Gln Phe Arg Tyr Ser Ser Thr Gly Glu Ile Ser Met Lys Leu Thr
 15 20 25

cca aaa gag tta gat aag ttg atg ctc cac tac gct gga gaa ttg gct 145
 Pro Lys Glu Leu Asp Lys Leu Met Leu His Tyr Ala Gly Glu Leu Ala
 30 35 40

aaa aaa cgc aaa gaa aaa ggc att aag ctt aac tat gta gaa gca gta 193
 Lys Lys Arg Lys Glu Lys Gly Ile Lys Leu Asn Tyr Val Glu Ala Val
 45 50 55 60

gct ttg att agt gcc cat att atg gaa gaa gcg aga gct ggt aaa aag 241
 Ala Leu Ile Ser Ala His Ile Met Glu Glu Ala Arg Ala Gly Lys Lys
 65 70 75

act gcg gct gaa ttg atg caa gaa ggg cgc act ctt tta aaa cca gat 289
 Thr Ala Ala Glu Leu Met Gln Glu Gly Arg Thr Leu Leu Lys Pro Asp
 80 85 90

gat gtg atg gat ggc gtg gca agc atg atc cat gaa gtg ggt att gaa	337
Asp Val Met Asp Gly Val Ala Ser Met Ile His Glu Val Gly Ile Glu	
95 100 105	
gcg atg ttt cct gat ggg act aaa ctc gta acc gtg cat acc cct att	385
Ala Met Phe Pro Asp Gly Thr Lys Leu Val Thr Val His Thr Pro Ile	
110 115 120	
gag gcc aat ggt aaa tta gtt cct ggt gag ttg ttc tta aaa aat gaa	433
Glu Ala Asn Gly Lys Leu Val Pro Gly Glu Leu Phe Leu Lys Asn Glu	
125 130 135 140	
gac atc act atc aac gaa ggc aaa aaa gcc gtt agc gtg aaa gtt aaa	481
Asp Ile Thr Ile Asn Glu Gly Lys Lys Ala Val Ser Val Lys Val Lys	
145 150 155	
aat gtt ggc gac aga ccg gtt caa atc ggc tca cac ttc cat ttc ttt	529
Asn Val Gly Asp Arg Pro Val Gln Ile Gly Ser His Phe His Phe Phe	
160 165 170	
gaa gtg aat aga tgc cta gac ttt gac aga gaa aaa act ttc ggt aaa	577
Glu Val Asn Arg Cys Leu Asp Phe Asp Arg Glu Lys Thr Phe Gly Lys	
175 180 185	
cgc tta gac att gcg agc ggg aca gcg gta aga ttt gag cct ggc gaa	625
Arg Leu Asp Ile Ala Ser Gly Thr Ala Val Arg Phe Glu Pro Gly Glu	
190 195 200	
gaa aaa tcc gta gaa ttg att gac att ggc ggt aac aga aga atc ttt	673
Glu Lys Ser Val Glu Leu Ile Asp Ile Gly Gly Asn Arg Arg Ile Phe	
205 210 215 220	
gga ttt aac gca ttg gtt gat aga caa gca gac aac gaa agc aaa aaa	721
Gly Phe Asn Ala Leu Val Asp Arg Gln Ala Asp Asn Glu Ser Lys Lys	
225 230 235	
att gct tta cac aga gct aaa gag cgt ggt ttt cat ggc gct aaa agc	769
Ile Ala Leu His Arg Ala Lys Glu Arg Gly Phe His Gly Ala Lys Ser	
240 245 250	
gat gac aac tat gta aaa aca att aag gag taa gaa atg aaa aag att	817
Asp Asp Asn Tyr Val Lys Thr Ile Lys Glu Glu Met Lys Lys Ile	
255 260 265	
agc aga aaa gaa tat gtt tct atg tat ggt cct act aca ggc gat aaa	865
Ser Arg Lys Glu Tyr Val Ser Met Tyr Gly Pro Thr Thr Gly Asp Lys	
270 275 280	
gtg aga ttg ggc gat aca gac ttg atc gct gaa gta gaa cat gac tac	913
Val Arg Leu Gly Asp Thr Asp Leu Ile Ala Glu Val Glu His Asp Tyr	

285	290	295	
acc att tat ggc gaa gag ctt aaa ttc ggt ggc ggt aaa acc cta aga			961
Thr Ile Tyr Gly Glu Leu Lys Phe Gly Gly Lys Thr Leu Arg			
300	305	310	315
gaa ggc atg agc caa tct aac aac cct agc aaa gaa gag ttg gat tta			1009
Glu Gly Met Ser Gln Ser Asn Asn Pro Ser Lys Glu Glu Leu Asp Leu			
320	325		330
att atc act aac gct tta atc gtg gat tac acc ggt att tat aaa gcg			1057
Ile Ile Thr Asn Ala Leu Ile Val Asp Tyr Thr Gly Ile Tyr Lys Ala			
335	340		345
gat att ggt att aaa gat ggc aaa atc gct ggc att ggt aaa ggc ggt			1105
Asp Ile Gly Ile Lys Asp Gly Lys Ile Ala Gly Ile Gly Lys Gly Gly			
350	355		360
aac aaa gac atg caa gat ggc gtt aaa aac aat ctt agc gta ggt cct			1153
Asn Lys Asp Met Gln Asp Gly Val Lys Asn Asn Leu Ser Val Gly Pro			
365	370		375
gct act gaa gcc tta gcc ggt gaa ggt ttg atc gta acg gct ggt ggt			1201
Ala Thr Glu Ala Leu Ala Gly Glu Gly Leu Ile Val Thr Ala Gly Gly			
380	385	390	395
att gac aca cac atc cac ttc att tca ccc caa caa atc cct aca gct			1249
Ile Asp Thr His Ile His Phe Ile Ser Pro Gln Gln Ile Pro Thr Ala			
400	405		410
ttt gca agc ggt gta aca acc atg att ggt ggt gga acc ggt cct gct			1297
Phe Ala Ser Gly Val Thr Thr Met Ile Gly Gly Gly Thr Gly Pro Ala			
415	420		425
gat ggc act aat gcg act act atc act cca ggc aga aga aat tta aaa			1345
Asp Gly Thr Asn Ala Thr Thr Ile Thr Pro Gly Arg Arg Asn Leu Lys			
430	435		440
tgg atg ctc aga gcg gct gaa gaa tat tct atg aat tta ggt ttc ttg			1393
Trp Met Leu Arg Ala Ala Glu Glu Tyr Ser Met Asn Leu Gly Phe Leu			
445	450		455
gct aaa ggt aac gct tct aac gat gcg agc tta gcc gat caa att gaa			1441
Ala Lys Gly Asn Ala Ser Asn Asp Ala Ser Leu Ala Asp Gln Ile Glu			
460	465	470	475
gcc ggt gcg att ggc ttt gca att cac gaa gac tgg ggc acc act cct			1489
Ala Gly Ala Ile Gly Phe Ala Ile His Glu Asp Trp Gly Thr Thr Pro			
480	485		490
tct gca atc aat cat gcg tta gat gtt gcg gac aaa tac gat gtg caa			1537

Ser	Ala	Ile	Asn	His	Ala	Leu	Asp	Val	Ala	Asp	Lys	Tyr	Asp	Val	Gln	
			495					500					505			
gtc	gct	atc	gcc	aca	gac	act	ttg	aat	gaa	gcc	ggc	tgt	gta	gaa	gac	1585
Val	Ala	Ile	Ala	Thr	Asp	Thr	Leu	Asn	Glu	Ala	Gly	Cys	Val	Glu	Asp	
			510				515					520				
act	atg	gct	gct	att	gct	gga	cgc	act	atg	cac	act	ttc	cac	act	gaa	1633
Thr	Met	Ala	Ala	Ile	Ala	Gly	Arg	Thr	Met	His	Thr	Phe	His	Thr	Glu	
			525				530					535				
ggc	gct	ggc	ggc	gga	cac	gct	cct	gat	att	att	aaa	gta	gcc	ggc	gaa	1681
Gly	Ala	Gly	Gly	Gly	His	Ala	Pro	Asp	Ile	Ile	Lys	Val	Ala	Gly	Glu	
			540				545				550				555	
cac	aac	att	ctt	ccc	gct	tcc	act	aac	ccc	acc	atc	cct	ttc	acc	gtg	1729
His	Asn	Ile	Leu	Pro	Ala	Ser	Thr	Asn	Pro	Thr	Ile	Pro	Phe	Thr	Val	
				560					565					570		
aat	aca	gaa	gca	gag	cac	atg	gac	atg	ctt	atg	gtg	tgc	cac	cac	ttg	1777
Asn	Thr	Glu	Ala	Glu	His	Met	Asp	Met	Leu	Met	Val	Cys	His	His	Leu	
				575					580					585		
gat	aaa	agc	att	aaa	gaa	gat	ggt	cag	ttc	gct	gat	tca	agg	atc	cgc	1825
Asp	Lys	Ser	Ile	Lys	Glu	Asp	Val	Gln	Phe	Ala	Asp	Ser	Arg	Ile	Arg	
			590					595				600				
cct	caa	acc	att	gcg	gct	gaa	gac	act	ttg	cat	gac	atg	ggg	att	ttc	1873
Pro	Gln	Thr	Ile	Ala	Ala	Glu	Asp	Thr	Leu	His	Asp	Met	Gly	Ile	Phe	
			605				610					615				
tca	atc	acc	agt	tct	gac	tct	caa	gcg	atg	ggc	cgt	gtg	ggc	gaa	gtt	1921
Ser	Ile	Thr	Ser	Ser	Asp	Ser	Gln	Ala	Met	Gly	Arg	Val	Gly	Glu	Val	
					625					630					635	
atc	act	aga	act	tgg	caa	aca	gct	gac	aaa	aac	aag	aaa	gaa	ttt	ggc	1969
Ile	Thr	Arg	Thr	Trp	Gln	Thr	Ala	Asp	Lys	Asn	Lys	Lys	Glu	Phe	Gly	
				640						645					650	
cgc	ttg	aaa	gaa	gaa	aaa	ggc	gat	aac	gac	aac	ttc	agg	atc	aaa	cgc	2017
Arg	Leu	Lys	Glu	Glu	Lys	Gly	Asp	Asn	Asp	Asn	Phe	Arg	Ile	Lys	Arg	
			655						660					665		
tac	ttg	tct	aaa	tac	acc	att	aac	cca	gcg	atc	gct	cat	ggg	att	agc	2065
Tyr	Leu	Ser	Lys	Tyr	Thr	Ile	Asn	Pro	Ala	Ile	Ala	His	Gly	Ile	Ser	
			670					675				680				
gag	tat	gta	ggc	tca	gta	gaa	gtg	ggc	aaa	gtg	gct	gac	ttg	gta	ttg	2113
Glu	Tyr	Val	Gly	Ser	Val	Glu	Val	Gly	Lys	Val	Ala	Asp	Leu	Val	Leu	
			685				690					695				

tggtggcgtgaaa	cccccac	atgatc	atc	aaa	ggc	2161				
Trp Ser Pro Ala Phe Phe Gly Val Lys Pro Asn Met Ile Ile Lys Gly										
700	705	710		715						
ggattgcgtttagc	caaatggcgat	gcgaacgcttct	atccct			2209				
Gly Phe Ile Ala Leu Ser Gln Met Gly Asp Ala Asn Ala Ser Ile Pro										
720	725		730							
accccaacccggtttat	tacagagaaatgttc	gctcatcat	ggtaaa			2257				
Thr Pro Gln Pro Val Tyr Tyr Arg Glu Met Phe Ala His His Gly Lys										
735	740		745							
gctaaa	tacgatgcaaac	atcactttt	gtgtctcaa	gcggcttat	gac	2305				
Ala Lys Tyr Asp Ala Asn Ile Thr Phe Val Ser Gln Ala Ala Tyr Asp										
750	755		760							
aaaggcattaaa	gaagaa	ttaggac	cttgaa	agacaa	gtgttgccg	gta 2353				
Lys Gly Ile Lys Glu Glu Leu Gly Leu Glu Arg Gln Val Leu Pro Val										
765	770		775							
aaa	aat	tgcaga	aatatc	actaaa	aaagac	atgatgcaa	ttcaac	gacact 2401		
Lys Asn Cys Arg Asn Ile Thr Lys Lys Asp Met Gln Phe Asn Asp Thr										
780	785		790		795					
accgctcac	attgaa	gtcaat	cctgaa	acttac	catgtgttc	gtgat 2449				
Thr Ala His Ile Glu Val Asn Pro Glu Thr Tyr His Val Phe Val Asp										
800	805		810							
ggc	aaa	gaagta	acttct	aaa	ccagcc	aat	aaa	gtgagc	ttggcg	caa 2497
Gly Lys Glu Val Thr Ser Lys Pro Ala Asn Lys Val Ser Leu Ala Gln										
815	820		825							
ctcttt	agcatt	ttctag	gat	ttt	ttaggagca	acgctc	cttaga	tcc 2545		
Leu Phe Ser Ile Phe Asp Phe Leu Gly Ala Thr Leu Leu Arg Ser										
830	835		840							
ccggga	attggg	gatccg	ctagcc	cgccta	atgagc	gggctt	tttttt 2593			
Pro Gly Ile Gly Asp Pro Leu Ala Arg Leu Met Ser Gly Leu Phe Phe										
845	850		855							
ctcggg	cagcgt	tgggtc	ctggcc	acgggt	gcgcat	gatcgt	gctcct 2641			
Leu Gly Gln Arg Trp Val Leu Ala Thr Gly Ala His Asp Arg Ala Pro										
860	865		870							
gtcgtt	gaggac	ccggct	aggctg	gcgggg	ttgcct	tactgg	ttagca 2689			
Val Val Glu Asp Pro Ala Arg Leu Ala Gly Leu Pro Tyr Trp Leu Ala										
875	880		885		890					
gaatga	atcacc	gatacg	cga	gcgaac	gtgaag	cga	ctgctg	ctgcaa 2737		
Glu Ile Thr Asp Thr Arg Ala Asn Val Lys Arg Leu Leu Leu Gln										
895	900		905							

aac gtc tgc gac ctg agc aac aac atg aat ggt ctt cgg ttt ccg tgt	2785
Asn Val Cys Asp Leu Ser Asn Asn Met Asn Gly Leu Arg Phe Pro Cys	
910 915 920	
ttc gta aag tct gga aac gcg gaa gtc agc gct ctt cgg ctt cct cgc	2833
Phe Val Lys Ser Gly Asn Ala Glu Val Ser Ala Leu Pro Leu Pro Arg	
925 930 935	
tca ctg act cgc tgc gct cgg tcg ttc ggc tgc ggc gag cgg tat cag	2881
Ser Leu Thr Arg Cys Ala Arg Ser Phe Gly Cys Gly Glu Arg Tyr Gln	
940 945 950	
ctc act caa agg cgg taa tac ggt tat cca cag aat cag ggg ata acg	2929
Leu Thr Gln Arg Arg Tyr Gly Tyr Pro Gln Asn Gln Gly Ile Thr	
955 960 965	
cag gaa aga aca tgt gag caa aag gcc agc aaa agg cca gga acc gta	2977
Gln Glu Arg Thr Cys Glu Gln Lys Ala Ser Lys Arg Pro Gly Thr Val	
970 975 980	
aaa agg ccg cgt tgc tgg cgt ttt tcc ata ggc tcc gcc ccc ctg acg	3025
Lys Arg Pro Arg Cys Trp Arg Phe Ser Ile Gly Ser Ala Pro Leu Thr	
985 990 995 1000	
agc atc aca aaa atc gac gct caa gtc aga ggt ggc gaa acc cga cag	3073
Ser Ile Thr Lys Ile Asp Ala Gln Val Arg Gly Gly Glu Thr Arg Gln	
1005 1010 1015	
gac tat aaa gat acc agg cgt ttc ccc ctg gaa gct ccc tcg tgc gct	3121
Asp Tyr Lys Asp Thr Arg Arg Phe Pro Leu Glu Ala Pro Ser Cys Ala	
1020 1025 1030	
ctc ctg ttc cga ccc tgc cgc tta ccg gat acc tgt ccg cct ttc tcc	3169
Leu Leu Phe Arg Pro Cys Arg Leu Pro Asp Thr Cys Pro Pro Phe Ser	
1035 1040 1045	
ctt cgg gaa gcg tgg cgc ttt ctc aat gct cac gct gta ggt atc tca	3217
Leu Arg Glu Ala Trp Arg Phe Leu Asn Ala His Ala Val Gly Ile Ser	
1050 1055 1060	
gtt cgg tgt agg tcg ttc gct cca agc tgg gct gtg tgc acg aac ccc	3265
Val Arg Cys Arg Ser Phe Ala Pro Ser Trp Ala Val Cys Thr Asn Pro	
1065 1070 1075 1080	
ccg ttc agc ccg acc gct gcg cct tat ccg gta act atc gtc ttg agt	3313
Pro Phe Ser Pro Thr Ala Ala Pro Tyr Pro Val Thr Ile Val Leu Ser	
1085 1090 1095	
cca acc cgg taa gac acg act tat cgc cac tgg cag cag cca ctg gta	3361
Pro Thr Arg Asp Thr Thr Tyr Arg His Trp Gln Gln Pro Leu Val	

1100	1105	1110	
aca gga tta gca gag cga ggt atg tag gcg gtg cta cag agt tct			3406
Thr Gly Leu Ala Glu Arg Gly Met	Ala Val Leu Gln Ser Ser		
1115	1120	1125	
tga agt ggt ggc cta act acg gct aca cta gaa gga cag tat ttg gta			3454
Ser Gly Gly Leu Thr Thr Ala Thr Leu Glu Gly Gln Tyr Leu Val			
1130	1135	1140	
tct gcg ctc tgc tga agc cag tta cct tcg gaa aaa gag ttg gta gct			3502
Ser Ala Leu Cys	Ser Gln Leu Pro Ser Glu Lys Glu Leu Val Ala		
1145	1150	1155	
ctt gat ccg gca aac aaa cca ccg ctg gta gcg gtg gtt ttt ttg ttt			3550
Leu Asp Pro Ala Asn Lys Pro Pro Leu Val Ala Val Val Phe Leu Phe			
1160	1165	1170	
gca agc agc aga tta cgc gca gaa aaa aag gat ctc aag aag atc ctt			3598
Ala Ser Ser Arg Leu Arg Ala Glu Lys Lys Asp Leu Lys Lys Ile Leu			
1175	1180	1185	
tga tct ttt cta cgg ggt ctg acg ctc agt gga acg aaa act cac gtt			3646
Ser Phe Leu Arg Gly Leu Thr Leu Ser Gly Thr Lys Thr His Val			
1190	1195	1200	
aag gga ttt tgg tca tga gat tat caa aaa gga tct tca cct aga tcc			3694
Lys Gly Phe Trp Ser	Asp Tyr Gln Lys Gly Ser Ser Pro Arg Ser		
1205	1210	1215	
ttt taa att aaa aat gaa gtt tta aat caa tct aaa gta tat atg agt			3742
Phe	Ile Lys Asn Glu Val Leu Asn Gln Ser Lys Val Tyr Met Ser		
1220	1225	1230	
aaa ctt ggt ctg aca gtt acc aat gct taa tca gtg agg cac cta tct			3790
Lys Leu Gly Leu Thr Val Thr Asn Ala	Ser Val Arg His Leu Ser		
1235	1240	1245	
cag cga tct gtc tat ttc gtt cat cca tag ttg cct gac tcc ccg tcg			3838
Gln Arg Ser Val Tyr Phe Val His Pro	Leu Pro Asp Ser Pro Ser		
1250	1255	1260	
tgt aga taa cta cga tac ggg agg gct tac cat ctg gcc cca gtg ctg			3886
Cys Arg	Leu Arg Tyr Gly Arg Ala Tyr His Leu Ala Pro Val Leu		
1265	1270	1275	
caa tgataccgcg agaccacgc tcaccggtc cagatttatc agcaataaac			3939
Gln			
cagccagccg gaagggccga gcgcagaagt ggtcctgcaa ctttatccgc ctccatccag			3999

tctattaatt gttgccggga agctagagta agtagttcgc cagttaatag tttgcgcaac 4059
 gttgttgcca ttgctgcagg catcgtggtg tcacgctcgt cgtttggtat ggcttcattc 4119
 agctccggtt cccaacgata aaggcgagtt acatgatecc ccatgttggtg caaaaaagcg 4179
 gttagctcct tcggtcctcc gatcgttggtc agaagtaagt tggccgcagt gttatcactc 4239
 atgggttatgg cagcactgca taattctctt actgtcatgc catccgtaag atgcttttct 4299
 gtgactggtg agtactcaac caagtcattc tgagaatagt gtatgcggcg accgagttgc 4359
 tcttgcccg cgtcaacacg ggataatacc gcgccacata gcagaacttt aaaagtgtc 4419
 atcattggaa aacgttcttc ggggcgaaaa ctctcaagga tcttaccgct gttgagatcc 4479
 agttcgtagt aaccctactc tgcacccaac tgatcttcag catcttttac tttcaccagc 4539
 gtttctgggt gagcaaaaac aggaaggcaa aatgccgcaa aaaagggaaat aaggcgaca 4599
 cggaaatgtt gaatactcat actcttctt tttcaatatt attgaagcat ttatcagggt 4659
 tattgtctca tgagcggata catatttgaa tgtatttaga aaaataaaca aataggggtt 4719
 ccgcgcacat ttccccgaaa agtgccacct gacgtctaag aaaccattat tatcatgaca 4779
 ttaacctata aaaataggcg tatcacgagg ccctttcgtc ttc aa 4824

<210> 2

<211> 10

<212> PRT

<213> Salmonella typhimurium

<400> 2

Asn Ser Ile Pro Glu Leu Arg Val Ile Lys

1

5

10

<210> 3

<211> 7

<212> PRT

<213> Salmonella typhimurium

<400> 3

Arg Thr Gln Gln Phe Arg Tyr

1

5

<210> 4

<211> 245

<212> PRT

<213> Artificial Sequence

<220>

<223> includes sequences from Salmonella typhimurium and
 Helicobacter pylori.

<400> 4

Ser Ser Thr Gly Glu Ile Ser Met Lys Leu Thr Pro Lys Glu Leu Asp

1

5

10

15

Lys Leu Met Leu His Tyr Ala Gly Glu Leu Ala Lys Lys Arg Lys Glu

20

25

30

Lys Gly Ile Lys Leu Asn Tyr Val Glu Ala Val Ala Leu Ile Ser Ala

35

40

45

His Ile Met Glu Glu Ala Arg Ala Gly Lys Lys Thr Ala Ala Glu Leu
 50 55 60
 Met Gln Glu Gly Arg Thr Leu Leu Lys Pro Asp Asp Val Met Asp Gly
 65 70 75 80
 Val Ala Ser Met Ile His Glu Val Gly Ile Glu Ala Met Phe Pro Asp
 85 90 95
 Gly Thr Lys Leu Val Thr Val His Thr Pro Ile Glu Ala Asn Gly Lys
 100 105 110
 Leu Val Pro Gly Glu Leu Phe Leu Lys Asn Glu Asp Ile Thr Ile Asn
 115 120 125
 Glu Gly Lys Lys Ala Val Ser Val Lys Val Lys Asn Val Gly Asp Arg
 130 135 140
 Pro Val Gln Ile Gly Ser His Phe His Phe Phe Glu Val Asn Arg Cys
 145 150 155 160
 Leu Asp Phe Asp Arg Glu Lys Thr Phe Gly Lys Arg Leu Asp Ile Ala
 165 170 175
 Ser Gly Thr Ala Val Arg Phe Glu Pro Gly Glu Glu Lys Ser Val Glu
 180 185 190
 Leu Ile Asp Ile Gly Gly Asn Arg Arg Ile Phe Gly Phe Asn Ala Leu
 195 200 205
 Val Asp Arg Gln Ala Asp Asn Glu Ser Lys Lys Ile Ala Leu His Arg
 210 215 220
 Ala Lys Glu Arg Gly Phe His Gly Ala Lys Ser Asp Asp Asn Tyr Val
 225 230 235 240
 Lys Thr Ile Lys Glu
 245

<210> 5

<211> 570

<212> PRT

<213> Helicobacter pylori

<400> 5

Glu Met Lys Lys Ile Ser Arg Lys Glu Tyr Val Ser Met Tyr Gly Pro
 1 5 10 15
 Thr Thr Gly Asp Lys Val Arg Leu Gly Asp Thr Asp Leu Ile Ala Glu
 20 25 30
 Val Glu His Asp Tyr Thr Ile Tyr Gly Glu Glu Leu Lys Phe Gly Gly
 35 40 45
 Gly Lys Thr Leu Arg Glu Gly Met Ser Gln Ser Asn Asn Pro Ser Lys
 50 55 60
 Glu Glu Leu Asp Leu Ile Ile Thr Asn Ala Leu Ile Val Asp Tyr Thr
 65 70 75 80
 Gly Ile Tyr Lys Ala Asp Ile Gly Ile Lys Asp Gly Lys Ile Ala Gly
 85 90 95
 Ile Gly Lys Gly Gly Asn Lys Asp Met Gln Asp Gly Val Lys Asn Asn
 100 105 110
 Leu Ser Val Gly Pro Ala Thr Glu Ala Leu Ala Gly Glu Gly Leu Ile
 115 120 125
 Val Thr Ala Gly Gly Ile Asp Thr His Ile His Phe Ile Ser Pro Gln

130	135	140
Gln Ile Pro Thr Ala Phe	Ala Ser Gly Val Thr	Thr Met Ile Gly Gly
145	150	155
Gly Thr Gly Pro Ala Asp	Gly Thr Asn Ala Thr	Thr Ile Thr Pro Gly
165	170	175
Arg Arg Asn Leu Lys Trp	Met Leu Arg Ala Ala	Glu Glu Tyr Ser Met
180	185	190
Asn Leu Gly Phe Leu Ala	Lys Gly Asn Ala Ser	Asn Asp Ala Ser Leu
195	200	205
Ala Asp Gln Ile Glu Ala	Gly Ala Ile Gly Phe	Ala Ile His Glu Asp
210	215	220
Trp Gly Thr Thr Pro Ser	Ala Ile Asn His Ala	Leu Asp Val Ala Asp
225	230	235
Lys Tyr Asp Val Gln Val	Ala Ile Ala Thr Asp	Thr Leu Asn Glu Ala
245	250	255
Gly Cys Val Glu Asp Thr	Met Ala Ala Ile Ala	Gly Arg Thr Met His
260	265	270
Thr Phe His Thr Glu Gly	Ala Gly Gly Gly His	Ala Pro Asp Ile Ile
275	280	285
Lys Val Ala Gly Glu His	Asn Ile Leu Pro Ala	Ser Thr Asn Pro Thr
290	295	300
Ile Pro Phe Thr Val Asn	Thr Glu Ala Glu His	Met Asp Met Leu Met
305	310	315
Val Cys His His Leu Asp	Lys Ser Ile Lys Glu	Asp Val Gln Phe Ala
325	330	335
Asp Ser Arg Ile Arg Pro	Gln Thr Ile Ala Ala	Glu Asp Thr Leu His
340	345	350
Asp Met Gly Ile Phe Ser	Ile Thr Ser Ser Asp	Ser Gln Ala Met Gly
355	360	365
Arg Val Gly Glu Val Ile	Thr Arg Thr Trp Gln	Thr Ala Asp Lys Asn
370	375	380
Lys Lys Glu Phe Gly Arg	Leu Lys Glu Glu Lys	Gly Asp Asn Asp Asn
385	390	395
Phe Arg Ile Lys Arg Tyr	Leu Ser Lys Tyr Thr	Ile Asn Pro Ala Ile
405	410	415
Ala His Gly Ile Ser Glu	Tyr Val Gly Ser Val	Glu Val Gly Lys Val
420	425	430
Ala Asp Leu Val Leu Trp	Ser Pro Ala Phe Phe	Gly Val Lys Pro Asn
435	440	445
Met Ile Ile Lys Gly Gly	Phe Ile Ala Leu Ser	Gln Met Gly Asp Ala
450	455	460
Asn Ala Ser Ile Pro Thr	Pro Gln Pro Val Tyr	Tyr Arg Glu Met Phe
465	470	475
Ala His His Gly Lys Ala	Lys Tyr Asp Ala Asn	Ile Thr Phe Val Ser
485	490	495
Gln Ala Ala Tyr Asp Lys	Gly Ile Lys Glu Glu	Leu Gly Leu Glu Arg
500	505	510
Gln Val Leu Pro Val Lys	Asn Cys Arg Asn Ile	Thr Lys Lys Asp Met
515	520	525
Gln Phe Asn Asp Thr Thr	Ala His Ile Glu Val	Asn Pro Glu Thr Tyr
530	535	540

His Val Phe Val Asp Gly Lys Glu Val Thr Ser Lys Pro Ala Asn Lys
 545 550 555 560
 Val Ser Leu Ala Gln Leu Phe Ser Ile Phe
 565 570

<210> 6
 <211> 59
 <212> PRT
 <213> Salmonella typhimurium

<400> 6
 Asp Phe Leu Gly Ala Thr Leu Leu Arg Ser Pro Gly Ile Gly Asp Pro
 1 5 10 15
 Leu Ala Arg Leu Met Ser Gly Leu Phe Phe Leu Gly Gln Arg Trp Val
 20 25 30
 Leu Ala Thr Gly Ala His Asp Arg Ala Pro Val Val Glu Asp Pro Ala
 35 40 45
 Arg Leu Ala Gly Leu Pro Tyr Trp Leu Ala Glu
 50 55

<210> 7
 <211> 67
 <212> PRT
 <213> Salmonella typhimurium

<400> 7
 Ile Thr Asp Thr Arg Ala Asn Val Lys Arg Leu Leu Leu Gln Asn Val
 1 5 10 15
 Cys Asp Leu Ser Asn Asn Met Asn Gly Leu Arg Phe Pro Cys Phe Val
 20 25 30
 Lys Ser Gly Asn Ala Glu Val Ser Ala Leu Pro Leu Pro Arg Ser Leu
 35 40 45
 Thr Arg Cys Ala Arg Ser Phe Gly Cys Gly Glu Arg Tyr Gln Leu Thr
 50 55 60
 Gln Arg Arg
 65

<210> 8
 <211> 141
 <212> PRT
 <213> Salmonella typhimurium

<400> 8
 Tyr Gly Tyr Pro Gln Asn Gln Gly Ile Thr Gln Glu Arg Thr Cys Glu
 1 5 10 15
 Gln Lys Ala Ser Lys Arg Pro Gly Thr Val Lys Arg Pro Arg Cys Trp
 20 25 30
 Arg Phe Ser Ile Gly Ser Ala Pro Leu Thr Ser Ile Thr Lys Ile Asp

35 40 45
 Ala Gln Val Arg Gly Gly Glu Thr Arg Gln Asp Tyr Lys Asp Thr Arg
 50 55 60
 Arg Phe Pro Leu Glu Ala Pro Ser Cys Ala Leu Leu Phe Arg Pro Cys
 65 70 75 80
 Arg Leu Pro Asp Thr Cys Pro Pro Phe Ser Leu Arg Glu Ala Trp Arg
 85 90 95
 Phe Leu Asn Ala His Ala Val Gly Ile Ser Val Arg Cys Arg Ser Phe
 100 105 110
 Ala Pro Ser Trp Ala Val Cys Thr Asn Pro Pro Phe Ser Pro Thr Ala
 115 120 125
 Ala Pro Tyr Pro Val Thr Ile Val Leu Ser Pro Thr Arg
 130 135 140

<210> 9
 <211> 20
 <212> PRT
 <213> Salmonella typhimurium

<400> 9
 Asp Thr Thr Tyr Arg His Trp Gln Gln Pro Leu Val Thr Gly Leu Ala
 1 5 10 15
 Glu Arg Gly Met
 20

<210> 10
 <211> 6
 <212> PRT
 <213> Salmonella typhimurium

<400> 10
 Ala Val Leu Gln Ser Ser
 1 5

<210> 11
 <211> 19
 <212> PRT
 <213> Salmonella typhimurium

<400> 11
 Ser Gly Gly Leu Thr Thr Ala Thr Leu Glu Gly Gln Tyr Leu Val Ser
 1 5 10 15
 Ala Leu Cys

<210> 12
 <211> 43

<212> PRT

<213> Salmonella typhimurium

<400> 12

Ser Gln Leu Pro Ser Glu Lys Glu Leu Val Ala Leu Asp Pro Ala Asn
 1 5 10 15
 Lys Pro Pro Leu Val Ala Val Val Phe Leu Phe Ala Ser Ser Arg Leu
 20 25 30
 Arg Ala Glu Lys Lys Asp Leu Lys Lys Ile Leu
 35 40

<210> 13

<211> 20

<212> PRT

<213> Salmonella typhimurium

<400> 13

Ser Phe Leu Arg Gly Leu Thr Leu Ser Gly Thr Lys Thr His Val Lys
 1 5 10 15
 Gly Phe Trp Ser
 20

<210> 14

<211> 11

<212> PRT

<213> Salmonella typhimurium

<400> 14

Asp Tyr Gln Lys Gly Ser Ser Pro Arg Ser Phe
 1 5 10

<210> 15

<211> 23

<212> PRT

<213> Salmonella typhimurium

<400> 15

Ile Lys Asn Glu Val Leu Asn Gln Ser Lys Val Tyr Met Ser Lys Leu
 1 5 10 15
 Gly Leu Thr Val Thr Asn Ala
 20

<210> 16

<211> 15

<212> PRT

<213> Salmonella typhimurium

<400> 16

Ser Val Arg His Leu Ser Gln Arg Ser Val Tyr Phe Val His Pro
1 5 10 15

<210> 17

<211> 8

<212> PRT

<213> Salmonella typhimurium

<400> 17

Leu Pro Asp Ser Pro Ser Cys Arg
1 5

<210> 18

<211> 14

<212> PRT

<213> Escherichia coli

<400> 18

Leu Arg Tyr Gly Arg Ala Tyr His Leu Ala Pro Val Leu Gln
1 5 10

<210> 19

<211> 4824

<212> DNA

<213> Artificial Sequence

<220>

<223> includes sequences from Helicobacter pylori,
Salmonella typhimurium, and Escherichia coli

<221> CDS

<222> (3893)...(3934)

<221> CDS

<222> (3938)...(4027)

<221> CDS

<222> (4031)...(4285)

<221> CDS

<222> (4289)...(4300)

<221> CDS

<222> (4304)...(4408)

<221> CDS

<222> (4412)...(4471)

<221> CDS
<222> (4475)...(4588)

<221> CDS
<222> (4592)...(4669)

<221> CDS
<222> (4673)...(4711)

<221> CDS
<222> (4715)...(4774)

<221> CDS
<222> (4784)...(4824)

<400> 19
gaattctatt ccggaacttc gcgttataaa atgaatctga cgtacacagc aatttagata 60
ttaatcatcc acaggagaga tctccatgaa actcacecca aaagagttag ataagttgat 120
gctccactac gctggagaat tggctaaaaa acgcaaagaa aaaggcatta agcttaacta 180
tgtagaagca gtagctttga ttagtgccca tattatggaa gaagcgagag ctggtaaaaa 240
gactgcggct gaattgatgc aagaaggcg cactctttta aaaccagatg atgtgatgga 300
tggcgtggca agcatgatcc atgaagtggg tattgaagcg atgtttcctg atgggactaa 360
actcgtaacc gtgcataccc ctattgaggg caatggtaaa ttagttcctg gtgagttgtt 420
cttaaaaaat gaagacatca ctatcaacga aggcataaaa gccgttagcg tgaaagttaa 480
aatggttggc gacagaccgg ttcaaatcgg ctccacttc catttctttg aagtgaatag 540
atgcctagac tttgacagag aaaaaacttt cggtaaagcg ttagacattg cgagcgggac 600
agcggtaaga tttgagcctg gcgaagaaaa atccgtagaa ttgattgaca ttggcggtaa 660
cagaagaatc tttgatttta acgcattggg tgatagacaa gcagacaacg aaagcaaaaa 720
aattgcttta cacagagcta aagagcgtgg ttttcatggc gctaaaagcg atgacaacta 780
tgtaaaaaca attaaggagt aagaaatgaa aaagattagc agaaaagaat atgtttctat 840
gtatggctct actacaggcg ataaagttag attgggcgat acagacttga tcgctgaagt 900
agaacatgac tacaccattt atggcgaga gcttaaatc ggtggcggta aaaccctaag 960
agaaggcatg agccaatcta acaaccctag caaagaagag ttggatttaa ttatcactaa 1020
cgctttaatc gtggattaca ccggtattta taaagcggat attgggtatta aagatggcaa 1080
aatcgctggc attggtaaaag gcggtaaaca agacatgcaa gatggcgtaa aaaacaatct 1140
tagcgtaggt cctgctactg aagccttagc cgggtgaagg ttgatcgtaa cggctgggtg 1200
tattgacaca cacatccact tcatttcacc ccaacaaatc cctacagctt ttgcaagcgg 1260
tgtaacaacc atgattggtg gtggaaccgg tcctgctgat ggcactaatg cgactactat 1320
cactccaggc agaagaaaatt taaaatggat gctcagagcg gctgaagaat attctatgaa 1380
tttaggtttc ttggctaaag gtaacgcttc taacgatgag agcttagccg atcaaattga 1440
agccggtgag attggctttg caattcaaga agactggggc accactcctt ctgcaatcaa 1500
tcatgcttta gatgttgagg acaaatacga tgtgcaagtc gctatcgcca cagacacttt 1560
gaatgaagcc ggttgtgtag aagacactat ggctgctatt gctggacgca ctatgcacac 1620
tttccacact gaaggcgctg gcggcgagca cgctcctgat attattaaag tagccggtga 1680
acacaacatt ctcccgctt ccactaacc caccatccct ttcaccgtga atacagaagc 1740
agagcacatg gacatgctta tgggtgtgcca ccactggat aaaagcatta aagaagatgt 1800
tcagttcgct gattcaagga tccgcectca aaccattgag gctgaagaca ctttgcata 1860
catggggatt ttctcaatca ccagttctga ctctcaagcg atgggcgctg tgggtgaagt 1920
tatcactaga acttggcaaa cagctgacaa aaacaagaaa gaatttggcc gcttgaaaga 1980
agaaaaaggc gataacgaca acttcaggat caaacgctac ttgtctaaat acaccattaa 2040
cccagcgatc gctcatggga ttagcgagta tgtaggttca gtgaagtggt gcaaagtggt 2100

tgacttggtta ttgtggagtc cagcattctt tggcgtgaaa cccaacatga tcatcaaagg 2160
 cggattcatt gcgttaagcc aaatgggcga tgcgaacgct tctatcccta cccacacaacc 2220
 gggtttattac agagaaatgt tcgctcatca tggtaaagct aaatacagatg caaacatcac 2280
 ttttgtgtct caagcggtt atgacaaagg cattaaagaa gaattaggac ttgaaagaca 2340
 agtgttgccg gtaaaaaatt gcagaaatat cactaaaaaa gacatgcaat tcaacgacac 2400
 taccgctcac attgaagtca atcctgaaac ttaccatgtg ttcgtggatg gcaaagaagt 2460
 aactttctaaa ccagccaata aagtgaagctt ggcgcaactc tttagcattt tctaggattt 2520
 tttaggagca acgctcctta gatccccggg aattggggat ccgctagccc gcctaattgag 2580
 cgggcttttt tttctcgggc agcgttgggt cctggccacg ggtgcgcag atcgtgctcc 2640
 tgcgttgtag gacccggtta ggctggcggg gttgccttac tggttagcag aatgaatcac 2700
 cgatacgcga gcgaacgtga agcgactgct gctgcaaaac gtctgcgacc tgagcaacaa 2760
 catgaatggt cttcgttttc cgtgtttcgt aaagtctgga aacgcggaag tcagcgtct 2820
 tccgcttct cgtcactga ctgctgcgc tggctcgttc ggctgcggcg agcggtatca 2880
 gctcactcaa aggcggtaat acggttatcc acagaatcag gggataacgc aggaaagaac 2940
 atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa aggcgcggtt gctggcggtt 3000
 ttccataggc tccgcccccc tgacgagcat cacaaaaatc gacgctcaag tcagagggtg 3060
 cgaaaccgga caggactata aagataccag gcgtttcccc ctggaagctc cctcgtgcgc 3120
 tctcctgttc cgacctgcc gcttaacgga tactgtccg cttttctccc ttcgggaagc 3180
 gtggcgcttt ctcaatgctc acgctgtagg tatctcagtt cgggtgtaggt cgttcgctcc 3240
 aagctgggct gtgtgcacga acccccgtt cagcccgacc gctgcgcctt atccggtaac 3300
 tatcgtcttg agtccaaccc ggtaagacac gacttatcgc cactggcagc agccactggt 3360
 aacaggatta gcagagcgag gtatgtaggc ggtgctacag agttcttgaa gtggtggcct 3420
 aactacggct acactagaag gacagtattt ggtatctgcg ctctgctgaa gccagttacc 3480
 ttcggaaaaa gagttggtag ctcttgatcc ggcaaaaaa ccaccgctgg tagcgggtggt 3540
 ttttttgttt gcaagcagca gattacgcgc agaaaaaaag gatctcaaga agatcctttg 3600
 atcttttcta cggggtctga cgtcagtggt aacgaaaact cacgttaagg gattttggtc 3660
 atgagattat caaaaaggat cttcacctag atccttttaa attaaaaatg aagttttaaa 3720
 tcaatctaaa gtatatatga gtaacttgg tctgacagtt accaatgctt aatcagtgag 3780
 gcacctatct cagcgatctg tctatttcgt tcatccatag ttgectgact ccccgctcgtg 3840
 tagataacta cgatacggga gggcttacca tctggcccca gtgctgcaat ga tac cgc 3898

Tyr Arg

1

gag acc cac gct cac cgg ctc cag att tat cag caa taa acc agc cag 3946
 Glu Thr His Ala His Arg Leu Gln Ile Tyr Gln Gln Thr Ser Gln
 5 10 15

ccg gaa ggg ccg agc gca gaa gtg gtc ctg caa ctt tat ccg cct cca 3994
 Pro Glu Gly Pro Ser Ala Glu Val Leu Gln Leu Tyr Pro Pro Pro
 20 25 30

tcc agt cta tta att gtt gcc ggg aag cta gag taa gta gtt cgc cag 4042
 Ser Ser Leu Leu Ile Val Ala Gly Lys Leu Glu Val Val Arg Gln
 35 40 45

tta ata gtt tgc gca acg ttg ttg cca ttg ctg cag gca tcg tgg tgt 4090
 Leu Ile Val Cys Ala Thr Leu Leu Pro Leu Leu Gln Ala Ser Trp Cys
 50 55 60

cac gct cgt cgt ttg gta tgg ctt cat tca gct ccg gtt ccc aac gat 4138
 His Ala Arg Arg Leu Val Trp Leu His Ser Ala Pro Val Pro Asn Asp

65	70	75	80	
caa ggc gag tta cat gat ccc cca tgt tgt gca aaa aag cgg tta gct				4186
Gln Gly Glu Leu His Asp Pro Pro Cys Cys Ala Lys Lys Arg Leu Ala				
	85	90	95	
cct tcg gtc ctc cga tcg ttg tca gaa gta agt tgg ccg cag tgt tat				4234
Pro Ser Val Leu Arg Ser Leu Ser Glu Val Ser Trp Pro Gln Cys Tyr				
	100	105	110	
cac tca tgg tta tgg cag cac tgc ata att ctc tta ctg tca tgc cat				4282
His Ser Trp Leu Trp Gln His Cys Ile Ile Leu Leu Leu Ser Cys His				
	115	120	125	
ccg taa gat gct ttt ctg tga ctg gtg agt act caa cca agt cat tct				4330
Pro Asp Ala Phe Leu Leu Val Ser Thr Gln Pro Ser His Ser				
	130	135	140	
gag aat agt gta tgc ggc gac cga gtt gct ctt gcc cgg cgt caa cac				4378
Glu Asn Ser Val Cys Gly Asp Arg Val Ala Leu Ala Arg Arg Gln His				
	145	150	155	
ggg ata ata ccg cgc cac ata gca gaa ctt taa aag tgc tca tca ttg				4426
Gly Ile Ile Pro Arg His Ile Ala Glu Leu Lys Cys Ser Ser Leu				
	160	165	170	
gaa aac gtt ctt ccg ggc gaa aac tct caa gga tct tac cgc tgt				4471
Glu Asn Val Leu Arg Gly Glu Asn Ser Gln Gly Ser Tyr Arg Cys				
	175	180	185	
tga gat cca gtt cga tgt aac cca ctc gtg cac cca act gat ctt cag				4519
Asp Pro Val Arg Cys Asn Pro Leu Val His Pro Thr Asp Leu Gln				
	190	195	200	
cat ctt tta ctt tca cca gcg ttt ctg ggt gag caa aaa cag gaa ggc				4567
His Leu Leu Leu Ser Pro Ala Phe Leu Gly Glu Gln Lys Gln Glu Gly				
	205	210	215	
aaa atg ccg caa aaa agg gaa taa ggg cga cac gga aat gtt gaa tac				4615
Lys Met Pro Gln Lys Arg Glu Gly Arg His Gly Asn Val Glu Tyr				
	220	225	230	
tca tac tct tcc ttt ttc aat att att gaa gca ttt atc agg gtt att				4663
Ser Tyr Ser Ser Phe Phe Asn Ile Ile Glu Ala Phe Ile Arg Val Ile				
	235	240	245	250
gtc tca tga gcg gat aca tat ttg aat gta ttt aga aaa ata aac aaa				4711
Val Ser Ala Asp Thr Tyr Leu Asn Val Phe Arg Lys Ile Asn Lys				
	255	260	265	
tag ggg ttc cgc gca cat ttc ccc gaa aag tgc cac ctg acg tct aag				4759

Gly Phe Arg Ala His Phe Pro Glu Lys Cys His Leu Thr Ser Lys
 270 275 280

aaa cca tta tta tca tgacattaa cct ata aaa ata ggc gta tca cga ggc 4810
 Lys Pro Leu Leu Ser Pro Ile Lys Ile Gly Val Ser Arg Gly
 285 290

cct ttc gtc ttc aa 4824
 Pro Phe Val Phe
 295

<210> 20
 <211> 14
 <212> PRT
 <213> Escherichia coli

<400> 20
 Tyr Arg Glu Thr His Ala His Arg Leu Gln Ile Tyr Gln Gln
 1 5 10

<210> 21
 <211> 30
 <212> PRT
 <213> Escherichia coli

<400> 21
 Thr Ser Gln Pro Glu Gly Pro Ser Ala Glu Val Val Leu Gln Leu Tyr
 1 5 10 15
 Pro Pro Pro Ser Ser Leu Leu Ile Val Ala Gly Lys Leu Glu
 20 25 30

<210> 22
 <211> 85
 <212> PRT
 <213> Escherichia coli

<400> 22
 Val Val Arg Gln Leu Ile Val Cys Ala Thr Leu Leu Pro Leu Leu Gln
 1 5 10 15
 Ala Ser Trp Cys His Ala Arg Arg Leu Val Trp Leu His Ser Ala Pro
 20 25 30
 Val Pro Asn Asp Gln Gly Glu Leu His Asp Pro Pro Cys Cys Ala Lys
 35 40 45
 Lys Arg Leu Ala Pro Ser Val Leu Arg Ser Leu Ser Glu Val Ser Trp
 50 55 60
 Pro Gln Cys Tyr His Ser Trp Leu Trp Gln His Cys Ile Ile Leu Leu
 65 70 75 80
 Leu Ser Cys His Pro

85

<210> 23
 <211> 4
 <212> PRT
 <213> Escherichia coli

<400> 23
 Asp Ala Phe Leu
 1

<210> 24
 <211> 35
 <212> PRT
 <213> Escherichia coli

<400> 24
 Leu Val Ser Thr Gln Pro Ser His Ser Glu Asn Ser Val Cys Gly Asp
 1 5 10 15
 Arg Val Ala Leu Ala Arg Arg Gln His Gly Ile Ile Pro Arg His Ile
 20 25 30
 Ala Glu Leu
 35

<210> 25
 <211> 20
 <212> PRT
 <213> Escherichia coli

<400> 25
 Lys Cys Ser Ser Leu Glu Asn Val Leu Arg Gly Glu Asn Ser Gln Gly
 1 5 10 15
 Ser Tyr Arg Cys
 20

<210> 26
 <211> 38
 <212> PRT
 <213> Escherichia coli

<400> 26
 Asp Pro Val Arg Cys Asn Pro Leu Val His Pro Thr Asp Leu Gln His
 1 5 10 15
 Leu Leu Leu Ser Pro Ala Phe Leu Gly Glu Gln Lys Gln Glu Gly Lys
 20 25 30
 Met Pro Gln Lys Arg Glu
 35

<210> 27
<211> 26
<212> PRT
<213> Escherichia coli

<400> 27
Gly Arg His Gly Asn Val Glu Tyr Ser Tyr Ser Ser Phe Phe Asn Ile
1 5 10 15
Ile Glu Ala Phe Ile Arg Val Ile Val Ser
20 25

<210> 28
<211> 13
<212> PRT
<213> Escherichia coli

<400> 28
Ala Asp Thr Tyr Leu Asn Val Phe Arg Lys Ile Asn Lys
1 5 10

<210> 29
<211> 20
<212> PRT
<213> Salmonella typhimurium

<400> 29
Gly Phe Arg Ala His Phe Pro Glu Lys Cys His Leu Thr Ser Lys Lys
1 5 10 15
Pro Leu Leu Ser
20

<210> 30
<211> 13
<212> PRT
<213> Salmonella typhimurium

<400> 30
Pro Ile Lys Ile Gly Val Ser Arg Gly Pro Phe Val Phe
1 5 10

<210> 31
<211> 31
<212> DNA
<213> Helicobacter pylori

<400> 31

tagggaattc tcatgaaact caccocaaaa g

31

<210> 32

<211> 22

<212> DNA

<213> Helicobacter pylori

<400> 32

gccaaacttag cttcctttcg gg

22

<210> 33

<211> 34

<212> DNA

<213> Helicobacter pylori

<400> 33

tctactgcag gatccaaaat gctaaagagt tgcg

34

<210> 34

<211> 21

<212> DNA

<213> Salmonella typhimurium

<400> 34

tcaaattgga ccccttgctg a

21

<210> 35

<211> 20

<212> DNA

<213> Salmonella typhimurium

<400> 35

tattccggaa cttcgcgta

20

<210> 36

<211> 23

<212> DNA

<213> Helicobacter pylori

<400> 36

tgtttcctga tgggactaaa ctc

23

<210> 37

<211> 22

<212> DNA

<213> Helicobacter pylori

<400> 37

accaggaact aatttaccat tg

22

<210> 38

<211> 21
<212> DNA
<213> Helicobacter pylori

<400> 38
ttgattgaca ttggcggtaa c 21

<210> 39
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 39
gttgtctgct tgtctatcaa cc 22

<210> 40
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 40
ggtggcggtta aaaccctaag ag 22

<210> 41
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 41
ctttgctagg gttgtagat tg 22

<210> 42
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 42
aatccctaca gcttttgcaa gc 22

<210> 43
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 43
gtgccatcag caggaccggt tc 22

<210> 44
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 44
atcgccacag acactttgaa tg 22

<210> 45
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 45
tagcagccat agtgtcttct ac 22

<210> 46
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 46
tgaagacact ttgcatgaca tg 22

<210> 47
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 47
tgagagtcag aactggtgat tg 22

<210> 48
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 48
catgatcatc aaaggcggat tc 22

<210> 49
<211> 23
<212> DNA
<213> Helicobacter pylori

<400> 49
gaagcggttcg catcgcccat ttg 23

<210> 50
<211> 22
<212> DNA
<213> Helicobacter pylori

<400> 50
tcgtggatgg caaagaagta ac 22

<210> 51
<211> 20
<212> DNA
<213> Helicobacter pylori

<400> 51
gcgccaagct cactttattg 20

<210> 52
<211> 22
<212> DNA
<213> Salmonella typhimurium

<400> 52
caacgacagg agcacgatca tg 22